



Italian Journal of Gynæcology & Obstetrics

March 2023 - Vol. 35 - N. 1 - Quarterly - ISSN 2385 - 0868

Nausea and vomiting in pregnancy: a Systematic Review of socio-demographic risk factors

Pinelopi Varela^{1,2,*}, Anna Deltsidou²

¹ General Hospital of Athens "Alexandra", Athens, Greece.

² Department of Midwifery, University of West Attica, Athens, Greece.

ARTICLE INFO

History

Received: 08 March 2022

Received in revised form: 03 May 2022

Accepted: 17 May 2022

Available online: 21 March 2023

DOI: 10.36129/jog.2022.38

Key words

Socio-demographic; nausea; pregnancy; risk factors; vomiting.

*Corresponding author: Pinelopi Varela, RM, MSc, PhD(c). General Hospital of Athens "Alexandra", Lourou 4-2, Athens 115 28, Greece.
Email: pinelopimid@yahoo.gr.
ORCID: 0000-0002-9419-0778.

ABSTRACT

Objective. Nausea and vomiting in pregnancy are two of the most prevalent symptoms and may have an unfavourable effect on a pregnant's daily life. The aim of this systematic review was to identify studies which assessed the risk of socio-demographic factors for nausea and vomiting in pregnancy.

Materials and Methods. This study was conducted according to the PRISMA guidelines, and the PubMed, Scopus, and Google Scholar databases were reviewed. The Newcastle-Ottawa Scale (NOS) for non-randomized studies for the assessment of the studies' quality was used. The socio-demographic risk factors reviewed were age, educational level, income, employment, marital status, alcohol intake and smoking.

Results. Finally, 718 articles were found, of which 13 matched the eligibility criteria. The studies' sample sizes varied from 116 to 81,486 participants. The findings showed that results are conflicting for the majority of the risk variables studied. The majority of the studies did not identify age, educational level and alcohol consumption as risk factors for nausea and vomiting in pregnancy. The relationship between income, employment and marital status, smoking and nausea and vomiting in pregnancy remains ambiguous because of the existence of conflicting conclusions.

Conclusions. The emergence of nausea and vomiting in pregnancy is influenced by socio-demographic variables. Obstetric health care providers should be conscious of several socio-demographic characteristics which may act as risk factors for the occurrence of nausea and vomiting in pregnancy and they should use a holistic and sensitive approach to identify pregnant women who might have greater risks to experience these symptoms.

INTRODUCTION

During pregnancy, nausea and vomiting (NVP) are two of the most prevalent symptoms, affecting 35-91% of pregnancies [1]. While, according to a recent systematic review, these percentages range to 50-

90% of all pregnant women [2]. Previous studies have also reported that the prevalence rate for nausea is 50-80% and for vomiting and retching is 50% [3]. NVP frequently starts at the beginning of the first trimester (6-8 weeks) and usually diminishes at 16-20 weeks of gestation [4]. The most severe form

of NVP, hyperemesis gravidarum, looks to be on the far side of the spectrum of NVP [5, 6] and affects about 0.3-3% of pregnancies [3]. Hyperemesis gravidarum is the most prevalent reason for hospitalization in the first trimester of pregnancy [6, 7] because of a severe weight loss and dehydration [6, 8].

The cause of NVP is uncertain and inadequately comprehended. Several hypotheses have been suggested and it is considered to involve a number of factors such as a hormonal stimulus, evolutionary adaptation, and psychological predisposition [2, 6, 9, 10]. According to a study, the levels of human chorionic gonadotropin and estrogen are responsible since they coincide with the peak of symptoms of nausea and vomiting [11, 12]. The severity of NVP is linked to a worse quality of life and negative consequences on numerous aspects of daily life, social and professional functioning [13, 14, 15] as well as to birth outcomes [11, 16]. Because of its frequency, it is possible to be underestimated by obstetric care providers, and to that may contribute a woman's perception of the severity of her symptoms. Early detection and treatment of NVP may contribute to the avoidance of the development of hyperemesis gravidarum [6] and of more serious problems, such as hospitalization [6, 17].

Although the cause of NVP is doubtful, risk factors for it have aroused the research interest of the scientific community. Definite answers are still being sought around the question of whether socio-demographic characteristics can act as risk factors for NVP and furthermore which of them can actually be considered as risk factors for the occurrence of it. The recognizing and studying NVP's socio-demographic risk factors can be an effective step in the identification of women at risk. Moreover, adequate knowledge of the socio-demographic risk factors for NVP is essential for the planning of effective strategies focusing on supporting these pregnant. Therefore, the present study aimed to systematically review and identify studies which assessed the risk of socio-demographic factors for NVP.

METHODS

The PRISMA statement's (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines were followed in conducting this systematic review [18].

Eligibility criteria

Only cohorts and case controls, peer-reviewed, in English-language published studies between the years 2000 and 2020 were considered, with no limitations on geographic area. We considered studies that described or reported any diagnostic criteria for NVP and with samples not having hyperemesis gravidarum. The primary or one of the outcomes of the studies had to be the assessment of risk socio-demographic factors for NVP (*i.e.*, age, education level, income, employment and marital status, alcohol intake and smoking). Studies were excluded if they were unrelated to the study's goal and if they were not cohorts or case controls studies.

Search strategy

Studies were identified by searching in PubMed, Scopus, and Google Scholar databases, and this search has been completed in December 2021. Searched terms and filters were used for each database. **Table 1** summarizes the databases search strategy.

Study selection

The titles and abstracts of possibly relevant studies were separately assessed by the two researchers (V.P., D.A.) according to eligibility and exclusion criteria. The full texts of selected studies were retrieved for additional assessment. Consensus was used to resolve disagreements among reviewers.

Data extraction

The two authors (V.P., D.A.) worked individually to extract data from relevant studies using a pre-made piloted extraction template. Each study's data extraction was comprised: first author, year of publication, location of study, type of study, sample size and results given in terms of frequencies percent, OR (unadjusted or adjusted), 95% CIs and P-values when accessible.

Quality assessment

The authors (V.P., D.A.) assessed the quality of the included articles by using Newcastle Ottawa Scale (NOS) for non-randomized studies [19]. The NOS checklist evaluates studies in three areas based on their quality: selection of the study groups; compa-

Table 1. Databases search strategy.

Database	Key words	Filters
PubMed	(nausea) AND (vomiting) AND (pregnancy) AND (risk factors)	Humans English Female Adult: +19 years From: 2000-2020
Scopus	nausea AND vomiting AND pregnancy AND risk AND factors	From: 2000-2020 English Publication Stage: final
Google Scholar	nausea AND vomiting AND pregnancy AND risk factors	2000-2020

rability of the groups; and ascertainment of either the exposure or outcome of interest for case-control or cohort studies, respectively. The NOS “star system” has a minimum score of 0 and a maximum score of 9. We gave the studies a star rating based on their quality; low ratings imply low quality and a significant risk of bias. Studies with a score of less than six stars were deemed low-quality, those with a score of six to seven stars were deemed moderate-quality, and those with a score of seven stars or more were deemed high-quality.

RESULTS

A systematic search resulted in the identification of 718 studies. After removing duplicates (n = 127), the remaining 591 studies were screened. After exclusions with reasons (*i.e.*, title, abstract, full-text, review articles, and not English-language papers), 13 full-text studies remained to be assessed and all of them were included in this systematic review. The Prisma flow diagram of the selection process is shown in **Figure 1**.




Figure 1. PRISMA flow diagram showing the selection process for the studies inclusion.

Table 2. Characteristics of the studies included in the systematic review of sociodemographic risk factors for NVP.

First Author, (Year), Country	Type of study	Sample size	Main results	NOS Quality score
Weigel <i>et al.</i> (2000), USA [20]	Cohort study	n = 1,000	<p>Age: Maternal age was not associated with the risk for developing NVP</p> <p>Education: NVP was more common in women who had < 6 years of education</p> <p>Nausea: < 6 years: Adj OR 1.32, 95%CI 1.03-1.60 ≥ 6 years: Adj OR 1</p> <p>Vomiting: < 6 years: Adj OR 1.47, 95%CI 1.20-1.74 ≥ 6 years: Adj OR 1</p> <p>Employment: Vomiting but not nausea was more frequent in women who were white collars or professionals</p> <p>Vomiting: white collars or professionals: Adj OR 1.55, 95%CI 1.15-1.95 Housewife: Adj OR 1 Blue collar: Adj OR 0.86, 95%CI 0.63-1.10</p> <p>Nausea: white collars or professionals: Adj OR 1.26, 95%CI 0.85-1.67 Housewife: Adj OR 1 Blue collar: Adj OR 0.96, 95%CI 0.73-1.21</p> <p>Marital status: NVP was more common in women who were involved in informal marriages</p> <p>Nausea: Common law union: Adj OR 1.31, 95%CI 1.09-1.53 Single: Adj OR 1 Legally married: Adj OR 0.84, 95%CI 0.62-1.06</p> <p>Vomiting: Common law union: Adj OR 1.36, 95%CI 1.15-1.57 Single: Adj OR 1 Legally married: Adj OR 0.81, 95%CI 0.59-1.02</p>	7 moderate
Kallen <i>et al.</i> (2003), Sweden [21]	Cohort study	n = 3,675 NVP group: n = 2,906 No NVP group: n = 769	<p>Age: The major increase in risk for NVP has been identified in women aged younger than 25 years</p> <p>age < 19: Adj OR 1.64, 95%CI 0.68-3.94 age 20-24: Adj OR 1.34, 95%CI 1.03-1.75 age 25-29: Adj OR 1.12, 95%CI 0.94-1.33 age 30-34: Adj OR 0.92, 95%CI 0.78-1.09 age 35-39: Adj OR 0.76, 95%CI 0.61-0.95 age 40 >: Adj OR 0.77, 95%CI 0.44-1.33</p> <p>Employment: Women who worked outside home had a lower risk for NVP than housewives/women on maternity leave or who were unemployed</p> <p>Working outside home: Adj OR 1.00 Housewife or unemployed: Adj OR 2.86, 95%CI 1.49-5.46</p> <p>Other/unknown (students and women who for reasons had leave of absence from work): Adj OR 2.02, 95%CI 0.79-5.14</p> <p>Smoking: Smokers before pregnancy had a lower risk for NVP</p> <p>No smoking: Adj OR 1.00 < 10 cig./day: Adj OR 0.78, 95%CI 0.60-1.00 +10 cig./day: Adj OR 0.70, 95%CI 0.55-0.88</p>	5 low
Lindseth & Vari (2005), USA [22]	Cohort study	n = 116	<p>Age: Women who reported NVP were older ($t = 2.4, p = 0.02$) than those not experiencing NVP</p> <p>Education: No difference between the NVP group and the non NVP group ($t = 0.47, p = ns$)</p> <p>Alcohol intake: Women with higher late NVP scores were found to be significantly more severe alcohol users ($r = 0.26, p = 0.04$) during their pregnancy</p> <p>Smoking: no significant difference between the groups</p>	4 low
Louik <i>et al.</i> (2006), USA [23]	Case-control study	NVP group: n = 14,998 No NVP group: n = 7,489	<p>Age: There was a strong inverse relationship between age and the risk of NVP. Compared with younger women, the adjusted OR for women 40 or older was lower (Adj OR 0.47, 95%CI 0.37-0.60)</p> <p>Income: No difference in the incidence of NVP</p> <p>< 10,000\$/year: Adj OR 1.03, 95%CI 0.88-1.21 vs 10,000-45 000\$/year: Adj OR 1.09, 95%CI 1.00 1.20</p> <p>Employment: No difference in the incidence of NVP</p> <p>Adj OR 1.05, 95%CI 0.98- 1.13</p> <p>Education: There was a statistically significant elevation in risk among more educated women, but the difference was not substantial</p> <p>12-15 years: Adj OR 1.09, 95%CI 1.01- 1.17 vs >15 years: Adj OR 1.11, 95%CI 1.03-1.21</p> <p>Smoking: Women who smoked during pregnancy were significantly less likely to experience NVP than those who discontinued smoking or had never smoked, during pregnancy: Adj OR 0.80, 95%CI 0.74-0.86 vs before pregnancy: Adj OR 1.06, 95%CI 0.98-1.13</p>	5 low
Markl <i>et al.</i> (2008), Germany [24]	Cohort study	n = 3,624	<p>Age: No association between age and NVP was found, Adj OR 0.97, 95%CI 0.94-0.99</p> <p>Marital status: Being single raised the risk of NVP, living with partner: Adj OR 1.00 living alone: Adj OR 1.49, 95%CI 1.24-1.79</p> <p>Smoking: Smokers were less likely to have NVP than non-smokers, smokers Adj OR 1 non-smokers Adj OR 2.03, 95%CI 1.02-4.05.</p>	7 moderate

First Author, (Year), Country	Type of study	Sample size	Main results	NOS Quality score
Lacasse <i>et al.</i> (2009), Canada [25]	Cohort study	n = 367 NVP group: n = 288 No NVP group: n = 79	Age: No association between age and NVP was found, Adj OR 0.95, 95%CI 0.88-1.02 Income: Women with lower household income were more likely to have NVP in the 1 st trimester of pregnancy Less than 40000 CDN\$/yr: Adj OR 1 Between 40000 & 79999 CDN\$/yr: Adj OR 0.35, 95%CI 0.13-0.93. Employment: No association between employment status and NVP was found Student/ not working: Adj OR 1 Working: Adj OR 0.56, 95%CI 0.22-1.42 Education: No association between education level and NVP was found University completed: Adj OR 1 University not completed: Adj OR 0.61, 95%CI 0.31-1.21 Marital status: Women living alone were more likely to have NVP in the 1 st trimester of pregnancy, but with no statistical significance With spouse or with someone (family or cotenant): Adj OR 1 Living alone: Adj OR 1.63, 95%CI 0.15-18.31 Alcohol intake: No association between alcohol and NVP was found, Use of alcohol before pregnancy: Adj OR 1.13, 95%CI 0.50-2.54 Use of alcohol during 1 st trimester: Adj OR 0.80, 95%CI 0.28-2.26 Smoking: Smokers during the 1 st trimester were less likely to have NVP, but with no statistical significance Smoking before pregnancy: Adj OR 1.07, 95%CI 0.37-3.05 Smoking during 1 st trimester: Adj OR 0.28, 95%CI 0.05-1.50	6 moderate
Chan <i>et al.</i> (2011), USA [26]	Cohort study	NVP group: n = 1,274 No NVP group: n = 279	Age: No association between age groups and NVP was found Age < 25: Adj OR 1.1, 95%CI 1.0-1.2 Age 25-29: Adj OR 1.0 Age 30-34: Adj OR 1.0, 95%CI 0.9-1.1 Age ≥ 35: Adj OR 0.8, 95%CI 0.7-0.9 Education: No association between education years and NVP was found < 12 years: Adj OR 1.0, 95%CI 0.9- 1.1 ≥12 to < 16 years: Adj OR 1.1, 95%CI 1.0-1.2 ≥ 16 years: Adj OR 1.0 Marital status: No association between marital status and NVP was found Married: Adj OR 1.0 Other: Adj OR 1.0, 95%CI 0.9-1.0 Alcohol intake: No association between alcohol use and NVP was found Any: Adj OR 0.9, 95%CI 0.7-1.1 None: Adj OR 1.0	6 moderate
Naumann <i>et al.</i> (2012), USA [27]	Cohort study	NVP group: n = 2,174 No NVP group: n = 557	Age: Younger maternal age was found to be an independent risk factor for NVP age 21-24: Adj OR 0.72, 95%CI 0.54- 0.97 age 25-29: Adj OR 0.73, 95%CI 0.54- 0.98 age 30-34: Adj OR 0.69, 95%CI 0.49-0.98 age 35-39: Adj OR 0.44, 95%CI 0.27- 0.72 age +40: Adj OR 0.18, 95%CI 0.05- 0.60 Alcohol intake: Any alcohol consumption during the 1 st trimester was not significant predictor of NVP (NVP group vs no NVP group, p = 0.06)	5 low
Kramer <i>et al.</i> (2013), Canada [32]	Cohort study	Early pregnancy: n = 551 Late pregnancy: n = 575	Employment: Not working women were twice as likely to report moderate and severe NVP in the early pregnancy Moderate NVP-not working: Adj OR 2.16, 95%CI 1.28-3.65, p = 0.004 Severe NVP- not working: Adj OR 2.06, 95%CI 1.08-3.93, p = 0.028 Smoking: NVP was less common or less severe at late pregnancy in women who smoked Moderate NVP-smoking: Adj OR 0.57, 95%CI 0.22-1.49, p = 0.252 Severe NVP- smoking: Adj OR 0.24, 95%CI 0.11-0.52, p ≤ 0.001	5 low
Temming <i>et al.</i> (2014), USA [28]	Cohort study	NVP group: n = 5,207 No NVP group: n = 76,279	Age: Women reporting NVP were more likely to be younger NVP group: Mean:26.8, SD: ± 5.1 No NVP group: Mean: 28.1, SD: ± 5.0, p ≤ 0.001 Marital status: Women reporting NVP were more likely to be single Unmarried: NVP group: 25.7% No NVP group: 19.0% OR 1.47, 95%CI 1.38- 1.57, p ≤ 0.001 Smoking: Women reporting NVP were more likely to smoke Smoking: NVP group: 3.3% No NVP group: 2.2% OR 1.51, 95%CI 1.29- 1.78, p ≤ 0.001.	6 moderate

First Author, (Year), Country	Type of study	Sample size	Main results	NOS Quality score
Choi <i>et al.</i> (2018), Korea [29]	Cohort study	n = 472 NVP group: n = 381 No NVP group: n = 91	Age: No significant differences were observed between the study groups < 35 age NVP group: n = 225 (59.1%) No NVP group: n = 61 (67.0%) ≥ 35 age NVP group: n = 156 (40.9%) No NVP group: n = 30 (33.0%), p = 0.160 Marital status: No significant differences were observed between the study groups Unmarried: NVP group: n = 4 (1.0%) No NVP group: n = 1 (1.1%) Married: NVP group: n = 377 (99.0%) No NVP group: n = 90 (98.9%), p = 1.000 Employment: No significant differences were observed between the study groups Unemployed: NVP group: n = 156 (40.9%) No NVP group: n = 43 (47.3%) Employed: NVP group: n = 225 (59.1%) No NVP group: n = 48 (52.7%), p = 0.270 Education: No significant differences were observed between the study groups Up to high school: NVP group: n = 34 (8.9%) No NVP group: n = 14 (15.4%) College/university or higher NVP group: n = 347 (91.1%) No NVP group: n = 77 (84.6%), p = 0.067 Alcohol intake: Alcohol consumption was associated with decreased odds of NVP No alcohol: Adj OR 1 Alcohol: Adj OR 0.36, 95%CI 0.14-0.93, p = 0.035 Smoking: Smoking was not associated with the NVP No smoking: Adj OR 1 Smoking: Adj OR 2.69, 95%CI 0.26-28.01, p = 0.410	5 low
Ellilä <i>et al.</i> (2018), Finland [30]	Cohort study	n = 2,381 moderate NVP: n = 1,243 mild NVP: n = 700 severe NVP: n = 152 no NVP: n = 286	Age: No association between age and NVP was found No NVP: OR 1 Mild NVP: OR 1, 95%CI 0.97-1.03 Moderate NVP: OR 0.99, 95%CI 0.96-1.02 Severe NVP: OR 0.97, 95%CI 0.93-1.01, p = 0.370 Marital status: No association between cohabited or single and NVP was found No NVP: OR 1 Mild NVP: OR 1.55, 95%CI 0.70-3.4 Moderate NVP: OR 0.81, 95%CI 0.41-1.61 Severe NVP: OR 0.59, 95%CI 0.24-1.49, p = 0.078 Employment: Working status (working/not working) was not associated with the severity of NVP No NVP: OR 1 Mild NVP: OR 1.19, 95%CI 0.79-1.78 Moderate NVP: OR 1.19, 95%CI 0.81-1.75 Severe NVP: OR 1.40, 95%CI 0.80-2.46, p = 0.684 Smoking: Smoking was not associated with the severity of NVP No NVP: OR 1 Mild NVP: OR 0.74, 95%CI 0.50-1.09 Moderate NVP: OR 0.74, 95%CI 0.51-1.06 Severe NVP: OR 0.79, 95%CI 0.45-1.39, p = 0.387	6 moderate
Regodón <i>et al.</i> (2020), Nepal [31]	Cohort study	1 st trimester: NVP group: n = 731 No NVP group: n = 475 2 nd trimester: NVP group: n = 1,137 No NVP group: n = 2,186 3 rd trimester: NVP group: n = 517 No NVP group: n = 2,884	Age: No association between mean age and NVP was found NVP group: Mean: 22.5, SD: 4.7 No NVP group: Mean: 22.6, SD: 4.7 Education: Overall it was not observed significant differences between the proportions of groups Literate: NVP group: n = 1,027 (59.7%) No NVP group: n = 1,076 (61.6%) No education: NVP group: n = 736 (42.8%) No NVP group: n = 738 (42.3%) When comparing women with symptoms vs without symptoms in the third trimester specifically, it was observed a statistically significant lower proportion of literate women in the symptomatic group (53.8% vs 62.4%, p ≤ 0.001) Smoking: Overall it was not observed significant differences between the proportions of groups NVP group: n = 70 (3.9%) No NVP group: n = 76 (4.2%)	4 low

Characteristics of the studies

Twelve studies had a cohort design, and one study had a case control design. The included studies were published between 2000 and 2020. Six studies were conducted in the USA, two in Canada, one in Germany, one in Sweden, one in Nepal, one in Finland and one in Korea. The sample size of the studies ranged between 116 and 81,486 participants. Six studies were of moderate quality and seven studies were of low quality. **Table 2** shows the main features and characteristics of the included studies.

Age

The connection between age and NVP was studied in twelve studies [20-31]. The majority of the studies (7/12) did not identify age as a risk factor for NVP [20, 24-26, 29-31]. Younger women were more likely to experience NVP in the studies of Louik *et al.* [23], Naumann *et al.* [27], and Temming *et al.* [28]. These findings are in agreement with Kallen's findings which showed that the major increase in risk occurred in women aged younger than 25 years [21]. Lindseth and Vari found that women who experienced NVP were older than those without NVP. However, the mean age of their sample was 26.3 years [22].

Income

Two studies examined the association between income and NVP with contradicting results. Louik *et al.* reported in their study that the incidence of NVP did not differ noticeably between groups [23]. In contrast, Lacasse *et al.* found that women having a lower family income (< 40,000 cdn\$/y) were more likely to have NVP in the first trimester of pregnancy than women with higher household income (between 40,000 and 79,999 cdn\$/y) [25].

Education

Seven studies investigated the link between educational level and NVP [20, 22, 23, 25, 26, 29, 31]. The level of education has not been found to be a risk factor for NVP in the studies of Lindseth and Vari [22], Louik *et al.* [23], Lacasse *et al.* [25], Chan *et al.* [26], Choi *et al.* [29], and Regodón *et al.* [31]. Louik *et al.* mentioned that though there was a statistically significant increase in risk among women having higher educational level, the difference was not

considerable, and they did not verify relationships between overall occurrence of NVP and low educational level [23]. In addition, Regodón *et al.* [31] reported that when comparing women with symptoms *vs* without symptoms in the third trimester, specifically, they observed a statistically significant lower proportion of literate women in the symptomatic group. However, generally, there were no significant variations in the percentages of the categories [31]. Nevertheless, the findings of the study of Weigel *et al.* do not support the previous results. In their study nausea and vomiting were more common in women who had fewer than six years of formal schooling [20].

Employment status

The relationship between employment status and NVP was studied in seven studies [20, 21, 23, 25, 29, 30, 32]. No association between employment status and NVP was found in the studies of Louik *et al.* [23], Lacasse *et al.* [25], and Choi *et al.* [29]. These results match those observed in the study of Ellilä *et al.* in which working status was not associated with the severity of NVP [30]. As opposed to the above findings are those of Kallen *et al.* who found that women who worked outside the home had a lower risk for NVP than housewives/women on maternity leave or who were not working [21]. Similarly, Kramer *et al.* stated that women who were unemployed were twice as likely to report moderate and severe NVP in the first/early second trimester of pregnancy [32]. Interestingly, Weigel *et al.* demonstrated that vomiting but not nausea was more frequent in women who were employees earning higher salaries in average doing highly skilled work (white collar or professionals) [20].

Marital status

Overall seven studies looked into the relationship between marital status and NVP. Two of the studies did not find any association between them [26, 29] and one study found that marital status was not associated with the severity of NVP [30]. Oppositely, Markl *et al.* observed that being single raised the risk of NVP, compared to women who lived with a partner [24]. Likewise, women reporting NVP were more likely to be unmarried in the study of Temming *et al.* [28]. In addition, Lacasse *et al.* stated that women living alone were more likely to have NVP in the first trimester of pregnancy, but there

was not a statistical significance [25]. Also, informal marriage retained its positive association with NVP according to the study of Weigel *et al.* [20].

Alcohol intake

Alcohol consumption as a possible risk factor for NVP, was examined by five studies. The majority of the studies (3/5) did not find any association between alcohol and NVP [25-27]. In the study of Choi *et al.* alcohol consumption was associated with decreased odds of NVP [29]. However, Lindseth and Vari demonstrated that women who experienced late NVP were found to have significantly more alcohol intakes during their pregnancy [22].

Smoking

On the whole ten studies examined the association between smoking and NVP. Three of them did not find any association between smoking and NVP [22, 29, 31] and in another one smoking was not associated with the severity of NVP [30]. Four studies [21, 23, 24, 32] suggested that smokers were less likely to experience NVP than non-smokers. These findings are consistent with those of Lacasse *et al.* who found that smokers during the first trimester were less likely to have NVP, but with no statistical significance [25]. In contrary to the above results Temming *et al.* found that women reporting NVP were more likely to smoke [28].

DISCUSSION

The purpose of this systematic review was to compile the evidence from observational studies and synthesize it regarding the socio-demographic risk factors of NVP. The current systematic review's results indicate that existing findings are contradictory for the most of the risk factors examined.

The results of the most studies have failed to identify the age as a risk factor for the occurrence of NVP. However, studies have shown that younger women are more likely to experience NVP. The existence of conflicting findings makes it difficult to reach an absolute conclusion. It is not easy to explain why younger ages were associated with NVP, but perhaps we can assume that younger women cope with pregnancy in a more stressful way. According to the literature, the hormones involved in the stress response (*e.g.*, cortisol) are positively related to the severity of nausea [33].

The findings of the current study cannot establish the generalizability of the association between income and symptoms of NVP, due to the fact that only two studies examined the above mentioned association, and furthermore, they used different definitions of household income categories (*i.e.*, cdn\$/yr and \$/yr). A comparative interpretation would be easier and clearer if the number of the studies were more extended, and the currency was the same. The educational level did not find to be a risk factor for NVP in the majority of the studies. Any possible pathway leading to a positive association remains unclear. It is apparent from the above results that there are contradicting findings relatively the association between employment status and NVP. Although not all studies have shown that unemployment can be a risk factor for NVP, the fact that it was found to be positively related in some of them cannot be ignored. It is not easy to interpret the significance of this association without knowing how many women were unable to work as a result of the NVP or this was of their own choice. This finding suggests that psychological variables may be important as well. The link between better health and consistent, high-quality work is well documented [34, 35]. It is not entirely definite whether marital status is a risk factor for NVP since there are findings that contradict each other. Single pregnant women may be more likely to experience nausea and vomiting, perhaps due to the existence of low social support or due to a stressful status affecting their health status [36]. For some pregnant, the absence of a partner may reflect financial and emotional insecurity which in turn it is possible to make them more fragile and vulnerable. Conflicting results exist regarding the link between alcohol consumption and NVP, thus making it difficult to determine whether alcohol is indeed a risk factor for NVP or not. Although the majority of the studies did not find any association between alcohol and NVP, there are studies with conflicting results. In one study alcohol use was linked with a lower risk of NVP [29], and in another one woman who was shown late NVP was found to use considerably higher alcohol amounts during her pregnancy [22]. Some authors have considered that alcohol consumers women compared to those of no consumers would be nausea or vomiting tolerant [29]. Another possible explanation for this is that alcohol intake lowers the hormone levels of estradiol and of human chorionic gonadotropin, which in turn lower the risk of

developing NVP [37]. Unknown factors may lead to the above inconsistency between the studies of Lindseth and Vari [22] and Choi *et al.* [29]. The existence of inconsistencies among studies' results regarding whether or not smoking is a risk factor for NVP, makes it complicated to find the answer. Interestingly, there are studies which demonstrated that smokers were less likely to experience NVP than non-smokers [21, 23, 24, 32]. These data support the protective effect of smoking against NVP and not that smoking may be a risk factor for NVP. Authors of previous studies have pointed out their suggestions on the subject. Kallen *et al.* hypothesized that "If NVP is an expression of a well-functioning placenta, the negative association with smoking may be because of a negative effect of maternal smoking on early placenta development" [21]. Also, the sense of odour [38] and taste [39] have been associated with increases in NVP and these senses may be decreased as a result of maternal smoking [32]. A variety of negative pregnancy outcomes have been associated with smoking during pregnancy, including placental abruption, miscarriage, stillbirth and preterm birth [40-42]. Health care professionals, and especially midwives, should encourage and support smoking cessation among all smoker pregnant women.

Interpretations regarding the presence of conflict findings might be described in part. The main reason can be considered to be the lack of homogeneity among the studies. Differences in methodological designs or the use of different NVP measures may have led to dissimilar conclusions. Furthermore, the samples' features and the confounding variables taken into account in the statistical analysis are also factors to consider.

One of the study's strengths is that it has been followed known recommendations (PRISMA) for systematic reviews of this type. In addition, it was used a reliable checklist for the assessment of the studies' quality. The present systematic review considered seven risk factors and the two reviewers worked separately based on predetermined criteria. The current review is limited by the lack of information in non-English papers and therefore some information might be missing. Another source of weakness that could have affected the findings is that solely three databases were used for the literature search.

Summarizing, the present review doubtless provides proof that socio-demographic factors do play

a role in the emergence of NVP. The majority of the studies did not identified age, level of education and alcohol consumption as a risk factor for NVP. The association between income, employment and marital status, smoking and NVP remains obscure since there are contradicting findings.

CONCLUSIONS

Nausea and vomiting in early pregnancy are common symptoms well-known to health care practitioners. Obstetricians, midwives and other health care providers should be conscious of several socio-demographic characteristics may act as risk factors for the occurrence of NVP. Therefore, they should be informed and up to date on the subject. Since there are conflicting research results more studies are needed. Future research will likely be able to bring to light answers by minimizing the divergences that already exist about socio-demographic risk factors for NVP. Furthermore, the findings of future research will add to the growing body of evidence indicating specific socio-demographic traits are indeed risk factors for NVP. More study data is needed by the scientific community in order to create supportive techniques for pregnant women who are more likely to experience NVP. Until further understanding and verification through scientific data on which socio-demographic factors and to what extent they are indeed risk factors for NVP, health professionals should use a holistic and sensitive approach to identify pregnant women who may have greater probabilities to experience these symptoms.

COMPLIANCE WITH ETHICAL STANDARDS

Authors contribution

P.V.: Conceptualization, data curation, formal analysis, investigation, methodology, project administration, writing – original draft. A.D.: Data curation, formal analysis, investigation, methodology, project administration, supervision, writing – review & editing.

Funding

None.

Study registration

N/A.

Disclosure of Interests

The authors declare that they have no conflict of interests.

Ethical approval

N/A.

Informed consent

N/A.

Data sharing

N/A.

REFERENCES

1. Einarson TR, Piwko C, Koren G. Quantifying the global rates of nausea and vomiting of pregnancy: a meta analysis. *J Popul Ther Clin Pharmacol.* 2013;20(2):e171-83. Available at: <https://pubmed.ncbi.nlm.nih.gov/23863575/>.
2. O'Donnell A, McParlin C, Robson SC, Beyer F, Moloney E, Bryant A, et al. Treatments for hyperemesis gravidarum and nausea and vomiting in pregnancy: a systematic review and economic assessment. *Health Technol Assess.* 2016;20(74):1-268. doi: 10.3310/hta20740.
3. Matthews A, Haas DM, O'Mathúna DP, Dowswell T. Interventions for nausea and vomiting in early pregnancy. *Cochrane database Syst Rev.* 2015;2015(9):CD007575. doi: 10.1002/14651858.CD007575.pub4.
4. Jarvis S, Nelson-Piercy C. Management of nausea and vomiting in pregnancy. *BMJ.* 2011;342(7812):1-8. doi: 10.1136/bmj.d3606.
5. Klebanoff MA, Koslowe PA, Kaslow R, Rhoads GG. Epidemiology of vomiting in early pregnancy. *Obs Gynecol.* 1985;66(5):612-6. Available at: <https://pubmed.ncbi.nlm.nih.gov/3903578/>.
6. Committee on Practice Bulletins-Obstetrics. ACOG Practice Bulletin No. 189: Nausea And Vomiting Of Pregnancy. *Obstet Gynecol.* 2018;131(1):e15-e30. doi: 10.1097/AOG.0000000000002456.
7. Gazmararian JA, Petersen R, Jamieson DJ, Schild L, Adams MM, Deshpande AD, et al. Hospitalizations during pregnancy among managed care enrollees. *Obstet Gynecol.* 2002;100(1):94-100. doi:10.1016/s0029-7844(02)02024-0.
8. Jueckstock JK, Kaestner R, Mylonas I. Managing hyperemesis gravidarum: a multimodal challenge. *BMC Med.* 2010;8:46. doi: 10.1186/1741-7015-8-46.
9. Simpson SW, Murphy Goodwin T, Robins SB, Rizzo AA, Howes RA, Buckwalter DK, et al. Psychological factors and hyperemesis gravidarum. *J Womens Health Gend Based Med.* 2001;10(5):471-7. doi: 10.1089/152460901300233948.
10. Flaxman SM, Sherman PW. Morning sickness: a mechanism for protecting mother and embryo. *Q Rev Biol.* 2000;75(2):113-48. doi: 10.1086/393377.
11. Furneaux EC, Langley-Evans AJ, Langley-Evans SC. Nausea and vomiting of pregnancy: endocrine basis and contribution to pregnancy outcome. *Obstet Gynecol Surv.* 2001;56(12):775-82. doi: 10.1097/00006254-200112000-00004.
12. McCarthy FP, Lutomski JE, Greene RA. Hyperemesis gravidarum: current perspectives. *Int J Womens Health.* 2014;6(1):719-25. doi: 10.2147/IJWH.S37685.
13. Lacasse A, Rey E, Ferreira E, Morin C, Bérard A. Nausea and vomiting of pregnancy: what about quality of life? *BJOG.* 2008;115(12):1484-93. doi: 10.1111/j.1471-0528.2008.01891.x.
14. Wood H, McKellar LV, Lightbody M. Nausea and vomiting in pregnancy: blooming or bloom-in' awful? A review of the literature. *Women Birth.* 2013;26(2):100-4. doi: 10.1016/j.wombi.2012.10.001.
15. Heitmann K, Nordeng H, Havnen GC, Solheimsnes A, Holst L. The burden of nausea and vomiting during pregnancy: severe impacts on quality of life, daily life functioning and willingness to become pregnant again - results from a cross-sectional study. *BMC Pregnancy Childbirth.* 2017;17(1):75. doi: 10.1186/s12884-017-1249-0.
16. Petry CJ, Ong KK, Beardsall K, Hughes IA, Acerini CL, Dunger DB. Vomiting in pregnancy is associated with a higher risk of low birth weight: a cohort study. *BMC Pregnancy Childbirth.* 2018;18(1):133. doi: 10.1186/s12884-018-1786-1
17. Brent R. Medical, social, and legal implications of treating nausea and vomiting of pregnancy. *Am J Obstet Gynecol.* 2002;186(5 Suppl Under-standing):S262-6. doi:10.1067/mob.2002.122603.

18. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Rev Esp Nutr Humana y Diet.* 2016;20(2):148-60. doi: 10.1186/2046-4053-4-1.
19. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for Assessing the Quality of Non-randomised Studies in Meta-Analyses. Ottawa Hosp Res Inst. 2019.
20. Weigel MM, Caiza ME, Lascano Y, Barreno G, Mosquera L. Early pregnancy nausea and vomiting in a high-altitude Andean population. *Int J Gynaecol Obstet.* 2000;69(1):9-21. doi: 10.1016/s0020-7292(99)00208-8.
21. Källén B, Lundberg G, Aberg A. Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. *Acta Obstet Gynecol Scand.* 2003;82(10):916-20. Available at: <https://pubmed.ncbi.nlm.nih.gov/12956841/>.
22. Lindseth G, Vari P. Nausea and vomiting in late pregnancy. *HealthCareWomenInt.* 2005;26(5):372-86. doi: 10.1080/07399330590933926.
23. Louik C, Hernandez-Diaz S, Werler MM, Mitchell AA. Nausea and vomiting in pregnancy: maternal characteristics and risk factors. *Paediatr Perinat Epidemiol.* 2006;20(4):270-8. doi: 10.1111/j.1365-3016.2006.00723.x.
24. Markl GE, Strunz-Lehner C, Egen-Lappe V, Lack N, Hasford J. The association of psychosocial factors with nausea and vomiting during pregnancy. *J Psychosom Obstet Gynaecol.* 2008;29(1):17-22. doi: 10.1080/01674820801902697.
25. Lacasse A, Rey E, Ferreira E, Morin C, Bérard A. Epidemiology of nausea and vomiting of pregnancy: prevalence, severity, determinants, and the importance of race/ethnicity. *BMC Pregnancy Childbirth.* 2009;9:26. doi: 10.1186/1471-2393-9-26.
26. Chan RL, Olshan AF, Savitz DA, Herring AH, Daniels JL, Peterson HB, et al. Maternal influences on nausea and vomiting in early pregnancy. *Matern Child Health J.* 2011;15(1):122-7. doi: 10.1007/s10995-009-0548-0.
27. Naumann CR, Zelig C, Napolitano PG, Ko CW. Nausea, vomiting, and heartburn in pregnancy: a prospective look at risk, treatment, and outcome. *J Matern Fetal Neonatal Med.* 2012;25(8):1488-93. doi: 10.3109/14767058.2011.644363.
28. Temming L, Franco A, Istwan N, Rhea D, Desch C, Stanziano G, et al. Adverse pregnancy outcomes in women with nausea and vomiting of pregnancy. *J Matern Fetal Neonatal Med.* 2014;27(1):84-8. doi: 10.3109/14767058.2013.806473.
29. Choi HJ, Bae YJ, Choi JS, Ahn HK, An HS, Hong DS, et al. Evaluation of nausea and vomiting in pregnancy using the Pregnancy-Unique Quantification of Emesis and Nausea scale in Korea. *Obstet Gynecol Sci.* 2018;61(1):30-7. doi: 10.5468/ogs.2018.61.1.30.
30. Ellilä P, Laitinen L, Nurmi M, Rautava P, Koivisto M, Polo-Kantola P. Nausea and vomiting of pregnancy: A study with pregnancy-unique quantification of emesis questionnaire. *Eur J Obstet Gynecol Reprod Biol.* 2018;230:60-7. doi: 10.1016/j.ejogrb.2018.09.031.
31. Regodón Wallin A, Tielsch JM, Khatry SK, Mullan LC, Englund JA, Chu H, et al. Nausea, vomiting and poor appetite during pregnancy and adverse birth outcomes in rural Nepal: an observational cohort study. *BMC Pregnancy Childbirth.* 2020;20(1):545. doi: 10.1186/s12884-020-03141-1.
32. Kramer J, Bowen A, Stewart N, Muhajarine N. Nausea and vomiting of pregnancy: prevalence, severity and relation to psychosocial health. *MCN Am J Matern Child Nurs.* 2013;38(1):21-7. doi: 10.1097/NMC.0b013e3182748489.
33. Otto B, Riepl RL, Klosterhalfen S, Enck P. Endocrine correlates of acute nausea and vomiting. *Auton Neurosci.* 2006;129(1-2):17-21. doi: 10.1016/j.autneu.2006.07.010.
34. Van Der Noordt M, IJzelenberg H, Droomers M, Proper KI. Health effects of employment: a systematic review of prospective studies. *Occup Environ Med.* 2014;71(10):730-6. doi: 10.1136/oemed-2013-101891.
35. Van Aerden K, Puig-Barrachina V, Bosmans K, Vanroelen C. How does employment quality relate to health and job satisfaction in Europe? A typological approach. *Soc Sci Med.* 2016;158:132-40. doi: 10.1016/j.socscimed.2016.04.017
36. Rodriguez A, Bohlin G, Lindmark G. Symptoms across pregnancy in relation to psychosocial and biomedical factors. *Acta Obstet Gynecol Scand.* 2001;80(3):213-23. doi: 10.1034/j.1600-0412.2001.080003213.x.
37. Weigel MM, Weigel RM. The association of reproductive history, demographic factors, and alcohol and tobacco consumption with the risk of developing nausea and vomiting in early pregnancy. *Am J Epidemiol.* 1988;127(3):562-70. doi: 10.1093/oxfordjournals.aje.a114831.

38. Swallow BL, Lindow SW, Masson EA, Hay DM. Women with nausea and vomiting in pregnancy demonstrate worse health and are adversely affected by odours. *J Obstet Gynaecol.* 2005;25(6):544-9. doi: 10.1080/01443610500230783.
39. O'Brien B, Naber S. Nausea and vomiting during pregnancy: effects on the quality of women's lives. *Birth.* 1992;19(3):138-43. doi: 10.1111/j.1523-536x.1992.tb00671.x.
40. Hammoud AO, Bujold E, Sorokin Y, Schild C, Krapp M, Baumann P, et al. Smoking in pregnancy revisited: findings from a large population-based study. *Am J Obstet Gynecol.* 2005;192(6):1856-62. doi: 10.1016/j.ajog.2004.12.057.
41. Salihu HM, Wilson RE. Epidemiology of prenatal smoking and perinatal outcomes. *Early Hum Dev.* 2007;83(11):713-20. doi: 10.1016/j.earlhumdev.2007.08.002.
42. Marufu TC, Ahankari A, Coleman T, Lewis S. Maternal smoking and the risk of still birth: systematic review and meta-analysis. *BMC Public Health.* 2015;15(1):239. doi: 10.1186/s12889-015-1552-5.