

## **NARRATIVE REVIEW**

### **Management of thin endometrium by hysteroscopic instillation of Platelet rich plasma: A review**

**Doi:** 10.36129/jog.2022.55

Rahul **Manchanda**<sup>1</sup>, Tarang **Yadav**<sup>2</sup>, Apoorva **Dave**<sup>3\*</sup>

#### **Authors' institutional affiliations:**

1. Department of gynae endoscopy, PSRI, New Delhi, India.
2. Fellow, Department of gynae endoscopy, PSRI, New Delhi, India.
3. Senior resident, Department of gynae endoscopy, PSRI, New Delhi, India.

\* **Corresponding author:** Apoorva **Dave**, Department of gynae endoscopy, PSRI, Saket, 110017 New Delhi, India. Email: dr.apoorvadave@gmail.com

#### **ORCID:**

Rahul **Manchanda** (0000-0002-6367-3005)

Tarang **Yadav** (0000-0002-7642-8500)

Apoorva **Dave** (0000-0001-8400-4241)

Word count: 1186

## Abstract

### **Management of thin endometrium by hysteroscopic instillation of Platelet rich plasma: A review**

**Introduction:** One of the prerequisites for successful pregnancy outcome is the endometrial thickness. In subfertile patients undergoing IVF treatment, the minimum endometrial thickness is presumed to be approximately 7 mm or more prior to embryo transfer. For the treatment of a refractory endometrium, instillation of platelet rich plasma in the subendometrial region via hysteroscopy is a newer technique. This study is conducted to evaluate the effectiveness of PRP treatment in cases of thin endometrium.

**Methods:** Review of various articles on hysteroscopic instillation of platelet rich plasma was undertaken by searching in the data bases like PubMed, Scopus, EMBASE, Web of Science, Science direct etc. The studies describing platelet rich plasma instillation and its different techniques including efficacy and their role in achieving clinical pregnancy were included in the present study. We searched studies from January 2000 to January 2022.

**Results:** Here in the previous studies we saw promising results with hysteroscopic Platelet rich plasma instillation in infertile women.

**Conclusion:** Autologous platelet rich plasma instillation is not associated with any side effects as it is derived from patients own blood. Also it is cost effective, less invasive, easily available as well as feasible for the specialist.

**Key words:** platelet rich plasma, hysteroscopy, PRP instillation

## **Management of thin endometrium by hysteroscopic instillation of Platelet rich plasma: A review**

### **Introduction and background:**

A functional embryo, a receptive endometrium and a synchronized interaction between blastocyst and endometrium are the prerequisite for the successful implantation [1]. Natural endometrial cycle comprises of three sequential phases Proliferative phase, secretory phase and menstrual phase [2]. During the 'window of implantation' phase that lasts for few days post-ovulation, the endometrium undergoes proliferation and increases in its thickness and becomes responsive to progesterone which is essential for the attachment, invasion and finally implantation of the embryo [2]. For embryo transfer the minimum endometrial thickness is presumed to be approximately 7 mm or more [3,4,5]. An important factor in implantation failure is considered to be a thin endometrium [6,7,8] due to high blood flow impedance of radial arteries of the uterine vessels, reduced epithelial growth, lower vascular development, and poor expression of vascular endothelial growth factor (VEGF) [9]. Several mechanisms have been proposed to explain the underlying pathophysiology of thin endometrium. Intrauterine adhesions, ovarian stimulation with clomiphene citrate (CC), as well as prolonged use of progesterone [10], or combined oral contraceptive pills have been associated with thin endometrium [11,12,13]

Several therapeutic approaches have been studied in the past which includes low-dose aspirin, pentoxifylline, tocopherol, L-arginine, vaginal sildenafil, extended estrogen, gonadotropin therapy, low-dose hCG, tamoxifen, acupuncture and neuromuscular electric stimulation, intrauterine G-CSF, and stem cell therapy [14]. A

derivative of fresh whole blood enriched with platelets autologous to plasma is known as Platelet rich plasma (PRP). PRP is considered to increase the receptivity of the endometrium via increasing vascularization, cell proliferation, anti-inflammatory properties and the reduced fibrosis, due to growth factors, peptides, and cytokines present in PRP [15,16]. For the treatment of a refractory endometrium, instillation of PRP in the subendometrial region via hysteroscopy is a newer technique. In this study we will review various articles on PRP instillation in cases of infertility having thin endometrium. The rationale behind this study is to understand the processes behind thin endometrium and focussing on its management with PRP which is a newer technique so that we can help women having subfertility and IVF failures due to this cause.

**Aims and objective:**

1. To study the effect of PRP instillation on endometrial thickness
2. To review the effect on pregnancy outcomes in patients undergoing PRP instillation

**Material and methods:**

A non-systematic review of various articles on hysteroscopic instillation of PRP was undertaken by searching in the data bases like PubMed, Scopus, EMBASE, Web of Science, Science direct etc. The studies describing PRP instillation and its different techniques including efficacy and their role in achieving clinical pregnancy were included in the present study. We searched studies from January 2000 to January 2022. The search strategy included keywords such as Platelet rich plasma, PRP, infertility, gynaecology, endometrium thickness, IVF, Recurrent implantation failure (RIF), hysteroscopic instillation, intrauterine infusion etc.

## **Discussion:**

In most of the studies the patient profile consists of history of repeated implantation failure (RIF), thin endometrium, intra uterine adhesion (IUA) [17] and chronic endometritis (CE), the PRP treatment is tried as no effective treatment is yet available for such cases [18-23]. It is shown that intrauterine PRP infusion gives successful reproductive outcomes, increases the thickness of endometrium and successful pregnancy outcomes thus can be used in various endometrial preparation protocols [18-23].

PRP increases the endometrial receptivity via increasing vascularity, cell proliferation and reduces fibrosis and inflammation due to various growth factors, peptides, and cytokines present in it [15,16].

PRP is made by taking blood from peripheral veins, it is rich in various growth factors like platelet-derived growth factor (PDGF), epidermal growth factor (EGF), transforming growth factor (TGF), VEGF, and cytokines which influences proliferation and growth of the endometrium [24].

Table 1 shows various studies done on PRP treatment in infertile cases and their outcomes. Zamaniyan M et al. conducted a study in Iran in a group of 98 women with RIF which showed positive clinical pregnancy outcomes with intrauterine infusion of PRP Significant higher clinical pregnancy rate in PRP infusion group (48.3%) as compared with control group (23%) [18].

Similar results were shown in RCT done by Nazari et al [19]. where Increased clinical and chemical pregnancy rates were observed in PRP group in comparison to control group (PRP infusion group vs control group: 45% vs 17% and 53% vs 27% respectively).

Mehrafza et al [20]. In their study done on group of 123 women and showed comparison between the outcome of PRP infusion with granulocyte colony stimulating factor(G-CSF) systemic injection and they showed significantly increased pregnancy outcomes with intrauterine infusion of PRP in contrast to systemic administration of G-CSF.

Eftekhar et al [21]. in their study on a group of 83 women, which included 40 cases and 43 controls and the case group were given PRP infusion and in the control group the embryo transfer was done without PRP infusion. The endometrial lining of the patients with PRP treatment increased significantly as compared to the controls ( $p=0.001$ ). The per-cycle clinical pregnancy rate was higher in the PRP group (32.5%) as compared to the control group (14%) but was not significant ( $p=0.044$ ) [14].

Zadehmodarres S et al. in their study on 10 patients infused 0.5 ml of PRP with IUI catheter showed 40% clinical pregnancy rate [22].

Another cross sectional study conducted by Agarwal et al. [23], where they injected 4ml PRP under hysteroscopic guidance in subendometrial junction using ovum pickup needle, among 32 primary and secondary infertility women showed improved ET in 24 patients and clinical pregnancy in 10 women and chemical pregnancy in 2 women.

The studies conducted by Wang X et al [25]., Tandulwadkar et al. [26] and Molina et al. [20] stated that the use of autologous PRP shows promising outcomes in the form of endometrium thickness and diverse pregnancy tests [25,26,27]. A meta-analysis done by Maleki et al. stated that the administered PRP at a dose range of 0.5–1 ml was more efficacious than those delivered PRP at the doses deviated from this range ( $\leq 0.5$  ml and  $\geq 1$  ml) [28].

Kaur et al. in their study on 98 patients with history of one previous FET failure with endometrium 7 mm or more in thickness and divided them into the control group and the intervention group (intrauterine infusion of 0.3-0.4ml of PRP). There was no increased pregnancy rate in the intervention group [29]. Therefore, the dose of the PRP used may have an important role to play in the outcomes.

**Conclusion:** The thin endometrium is a nightmare for an ART specialist as it poses challenges in embryo transfer and successful pregnancy outcome. Here in the previous studies we saw promising results with hysteroscopic PRP instillation in infertile women. Autologous PRP instillation is not associated with any side effects as PRP is derived from patients own blood. Also it is cost effective, less invasive, easily available as well as feasible for the specialist. Further randomised trials are needed for creating guidelines for the clinicians around the world. Additionally, we need standardised PRP preparation technique and also the precise indication of the PRP treatment needs to be published for its utility.

The authors report there are no competing interests to declare.

Compliance with Ethical standards: N/A

**Authors contribution:**

R.M.: Supervision

T.Y.: Conceptualization

A.D.: Writing- original draft, Writing- review & editing

No **funding** was received for this study.

**Ethical approval:** Not applicable

**Informed consent:** Not applicable

## References:

1. Craciunas L, Gallos I, Chu J, Bourne T, Quenby S, Brosens JJ, Coomarasamy A. Conventional and modern markers of endometrial receptivity: a systematic review and meta-analysis. *Hum Reprod Update*. 2019 Mar 1;25(2):202-223. doi: 10.1093/humupd/dmy044.
2. Simon A, Laufer N. Assessment and treatment of repeated implantation failure (RIF). *J Assist Reprod Genet*. 2012 Nov;29(11):1227-39. doi: 10.1007/s10815-012-9861-4. Epub 2012 Sep 14. PMID: 22976427; PMCID: PMC3510376.
3. El-Toukhy T, Coomarasamy A, Khairy M, Sunkara K, Seed P, Khalaf Y, Braude P. The relationship between endometrial thickness and outcome of medicated frozen embryo replacement cycles. *Fertil Steril*. 2008 Apr;89(4):832-9. doi: 10.1016/j.fertnstert.2007.04.031.
4. Richter KS, Bugge KR, Bromer JG, Levy MJ. Relationship between endometrial thickness and embryo implantation, based on 1,294 cycles of in vitro fertilization with transfer of two blastocyst-stage embryos. *Fertil Steril*. 2007 Jan;87(1):53-9. doi: 10.1016/j.fertnstert.2006.05.064.
5. Zadehmodarres S, Salehpour S, Saharkhiz N, Nazari L. Treatment of thin endometrium with autologous platelet-rich plasma: a pilot study. *JBRA Assist Reprod*. 2017 Feb 1;21(1):54-56. doi: 10.5935/1518-0557.20170013.
6. Abdalla HI, Brooks AA, Johnson MR, Kirkland A, Thomas A, Studd JW. Endometrial thickness: a predictor of implantation in ovum recipients? *Hum Reprod*. 1994 Feb;9(2):363-5. doi: 10.1093/oxfordjournals.humrep.a138509.

7. Alam V., Bernardini L., Gonzales J., Asch R.H., Balmaceda J.P. A prospective study of echographic endometrial characteristics and pregnancy rates during hormonal replacement cycles. *J. Assist. Reprod. Genet.* 1993;10:215–219. doi: 10.1007/BF01239224.
8. Gonen Y, Casper RF, Jacobson W, Blankier J. Endometrial thickness and growth during ovarian stimulation: a possible predictor of implantation in in vitro fertilization. *Fertil Steril.* 1989 Sep;52(3):446-50. doi: 10.1016/s0015-0282(16)60916-0.
9. Miwa I, Tamura H, Takasaki A, Yamagata Y, Shimamura K, Sugino N. Pathophysiologic features of "thin" endometrium. *Fertil Steril.* 2009 Apr;91(4):998-1004. doi: 10.1016/j.fertnstert.2008.01.029.
10. Svetlana Mateva, Margarita Nikolova, Kristin Yaneva, Angel Yordanov. Serous endometrial intraepithelial carcinoma coexistence with atypical polypoid adenomyoma in a young woman: a rare case. *Ital J Gynaecol Obstet.* 2022 June;34(2):71-75 doi: 10.36129/jog.2021.10.
11. H. Al-Inany Intrauterine adhesions. An update *Acta Obs Gynecol Scand*, 80 (11) (2001), pp. 986-993
12. Y. Gonen, R.F. Casper Sonographic determination of a possible adverse effect of clomiphene citrate on endometrial growth *Hum Reprod*, 5 (6) (1990), pp. 670-674
13. R.F. Casper It's time to pay attention to the endometrium *Fertil Steril*, 96 (3) (2011), pp. 519-521
14. Lebovitz O, Orvieto R. Treating patients with "thin" endometrium - an ongoing challenge. *Gynecol Endocrinol.* 2014 Jun;30(6):409-14. doi: 10.3109/09513590.2014.906571. Epub 2014 Apr 2. PMID: 24693854.

15. Bos-Mikich A, Ferreira MO, de Oliveira R, Frantz N. Platelet-Rich Plasma or Blood-Derived Products to Improve Endometrial Receptivity? *J Assist Reprod Genet* (2019) 36(4):613–20. doi: 10.1007/s10815-018-1386-z
16. Drago L, Bortolin M, Vassena C, Taschieri S, Del Fabbro M. Antimicrobial Activity of Pure Platelet-Rich Plasma Against Microorganisms Isolated From Oral Cavity. *BMC Microbiol* (2013) 13:47. doi: 10.1186/1471-2180-13-47
17. S. Jha, K. Surabhi Hysteroscopy “As one stop approach” in the management of intrauterine pathology. Focus on patient’s satisfaction. *Ital J Gynaecol Obstet*. 2021;33(2):102-109. doi: 10.36129/jog.2022.43.
18. Zamaniyan M, Peyvandi S, Heidaryan Gorji H, Moradi S, Jamal J, Yahya Poor Aghmashhadi F, Hossein Mohammadi M. Effect of platelet-rich plasma on pregnancy outcomes in infertile women with recurrent implantation failure: a randomized controlled trial. *Gynecol Endocrinol*. 2021 Feb;37(2):141-145. doi: 10.1080/09513590.2020.1756247.
19. Nazari L, Salehpour S, Hosseini MS, Hashemi Moghanjoughi P. The effects of autologous platelet-rich plasma in repeated implantation failure: a randomized controlled trial. *Hum Fertil (Camb)*. 2020 Sep;23(3):209-213. doi: 10.1080/14647273.2019.1569268.
20. Mehrafza M, Kabodmehri R, Nikpouri Z, Pourseify G, Raoufi A, Eftekhari A, Samadnia S, Hosseini A. Comparing the Impact of Autologous Platelet-rich Plasma and Granulocyte Colony Stimulating Factor on Pregnancy Outcome in Patients with Repeated Implantation Failure. *J Reprod Infertil*. 2019 Jan-Mar;20(1):35-41.

21. Eftekhari M, Neghab N, Naghshineh E, Khani P. Can autologous platelet rich plasma expand endometrial thickness and improve pregnancy rate during frozen-thawed embryo transfer cycle? A randomized clinical trial. *Taiwan J Obstet Gynecol.* 2018 Dec;57(6):810-813. doi: 10.1016/j.tjog.2018.10.007.
22. Zadehmodarres S, Salehpour S, Saharkhiz N, Nazari L. Treatment of thin endometrium with autologous platelet-rich plasma: a pilot study. *JBRA Assist Reprod.* 2017;21(1):54-56. Published 2017 Feb 1. doi:10.5935/1518-0557.20170013.
23. Agarwal M, Mettler L, Jain S, et al. Management of a thin endometrium by hysteroscopic instillation of platelet-rich plasma into the endomyometrial junction: a pilot study. *J Clin Med* 2020;9(9):2795. DOI: 10.3390/jcm9092795.
24. Chang Y, Li J, Chen Y, Wei L, Yang X, Shi Y, Liang X. Autologous platelet-rich plasma promotes endometrial growth and improves pregnancy outcome during in vitro fertilization. *Int J Clin Exp Med.* 2015 Jan 15;8(1):1286-90.
25. Wang X, Liu L, Mou S, Zhao H, Fang J, Xiang Y, Zhao T, Sha T, Ding J, Hao C. Investigation of platelet-rich plasma in increasing proliferation and migration of endometrial mesenchymal stem cells and improving pregnancy outcome of patients with thin endometrium. *J Cell Biochem.* 2018 Dec 3. doi: 10.1002/jcb.28014.
26. Tandulwadkar SR, Naralkar MV, Surana AD, Selvakarthick M, Kharat AH. Autologous Intrauterine Platelet-Rich Plasma Instillation for Suboptimal Endometrium in Frozen Embryo Transfer Cycles: A Pilot

Study. *J Hum Reprod Sci.* 2017 Jul-Sep;10(3):208-212. doi:

10.4103/jhrs.JHRS\_28\_17.

27. Molina A, Sánchez J, Sánchez W, Vielma V. Platelet-rich plasma as an adjuvant in the endometrial preparation of patients with refractory endometrium. *JBRA Assist Reprod.* 2018 Mar 1;22(1):42-48. doi: 10.5935/1518-0557.20180009.
28. Maleki-Hajiagha A, Razavi M, Rouholamin S, Rezaeinejad M, Maroufizadeh S, Sepidarkish M. Intrauterine infusion of autologous platelet-rich plasma in women undergoing assisted reproduction: A systematic review and meta-analysis. *J Reprod Immunol.* 2020 Feb;137:103078. doi: 10.1016/j.jri.2019.103078.
29. Kaur J, Madhavan A, Naidu P, Rani K, Mahajan N. Intrauterine Autologous Platelet-Rich Plasma Therapy to Improve Implantation Rates in Patients Undergoing Frozen Embryo Transfer: A Pilot Study. *Onco Fertil J.* 2018; 1(2):81–85. doi: 10.4103/tofj.tofj\_11\_18.

**Table1: Various studies on PRP instillation and their outcomes.**

| Study                       | Type of study              | No. of patients  | Method  | Outcome  |
|-----------------------------|----------------------------|--|---|--|
| Zamaniyan M et al.[18]      | Case control study         | 98 women with RIF                                      | 0.5ml intrauterine PRP infusion 48 hours prior to embryo transfer   | Significant higher clinical pregnancy rate in PRP infusion group(48.3%)                  |
| Nazari et al. [19]          | RCT                        | 138  | intrauterine PRP infusion group vs control group  | Increased clinical and chemical pregnancy rate in PRP group compared to controls         |
| Mehrafza et al. [20]        | Retrospective cohort study | 123  | compared the effect of intrauterine infusion of PRP with systemic administration of granulocyte colony stimulating factor(G-CSF)    | PRP more effective as compared to GCSF in RIF patients                                   |
| Eftekhari et al. [21]       | RCT                        | 83 (40 patient, 43 control)                            | 0.5-1.0ml intrauterine PRP infusion on the 13 <sup>th</sup> day of HRT cycle  | ET increased in patients as compared to control<br>Clinical pregnancy : same in 2 groups |
| Zadehmodarres S et al. [22] | Prospective cohort study   | 10   | Intrauterine infusion of 0.5 ml of PRP with the IUI catheter  | Clinical pregnancy: 4  |
| Agarwal et al.[23]          | Cross sectional            | 32 (secondary infertility:8, Primary infertility : 24) | Under hysteroscopic guidance 4 mL of PRP was injected into the subendometrial region using ovum pickup needle (1.0 mL in each wall) | Improved ET 24 patients<br>Clinical pregnancy: 10<br>Chemical pregnancy: 2               |