

CASE REPORT

A clitoral epidermal inclusion cyst in a postmenopausal woman: a case report

Short title: Clitoral cyst in adult population.

Federica **Savasta**^{1,*}, Alessandro **Libretti**¹, Alberto **Catinella**¹, Antonio **Taveggia**², Libera **Troia**¹, Valentino **Remorgida**¹

¹ Department of Gynecology and Obstetrics, University of Piemonte Orientale, Ospedale Maggiore della Carità, Novara, Italy.

² Department of Plastic Surgery, University of Piemonte Orientale, Ospedale Maggiore della Carità, Novara, Italy.

***Corresponding author:** Federica **Savasta**, Department of Gynaecology and Obstetrics, University of Piemonte Orientale, Ospedale Maggiore della Carità, corso Mazzini 19, Novara, 28100, Italy.

Email: federica_savasta@hotmail.it.

ORCID: 0009-0008-7722-7832.

Doi: 10.36129/jog.2024.152

ABSTRACT

Background. Clitoral cysts are rare conditions typically observed in the paediatric population, manifesting as either benign or malign entities, and may be congenital or acquired. The most prevalent forms are epidermal inclusion cysts, often associated with trauma, particularly female genital mutilation, although their aetiology can sometimes remain unclear.

This study aims to present an atypical case of a clitoral cyst occurring in an adult, post-menopausal woman, without any evident risk factor.

Case presentation. A 56-year-old post-menopause patient was referred to the emergency department with a 4 cm spontaneous swelling in the clitoral area. The patient denied any history of genital trauma and reported a gradual increase in size over the past two years, accompanied by discomfort and dyspareunia.

Upon clinical examination, ultrasound revealed a unilocular, hypoechoic, non-vascularized cystic structure. Subsequently, surgical excision was performed and pathologic examination confirmed it as an epidermal inclusion cyst.

To date, the literature reports only 14 cases of spontaneous clitoral epidermal inclusion cyst in post-puberal and adult patients, highlighting their exceedingly rare occurrence in adult women without a history of trauma.

Conclusions. Despite their benign and rare nature, epidermal inclusion clitoral cysts may induce significant discomfort, potentially leading to sexual dysfunction if left untreated. Therefore, it is imperative for clinicians to recognize them, considering the like hood of their occurrence and growth even post-menopausal women. Addressing this condition through proper counselling and prompt surgical intervention is crucial to ensure a satisfactory quality of life for affected patients.

Key words

Clitoral cyst; anorgasmia; dyspareunia; epidermal cyst; case report.

INTRODUCTION

Epidermal inclusion cysts are benign, spontaneous, cutaneous cysts that are usually found on the face, neck, back or scalp, and less commonly on the vulvar area, though they can occur in various locations. They are caused by the implantation of epidermal elements into the dermis layer of the skin and are usually filled with laminated keratinous material [1]. Epidermoid cysts of the clitoris are a very rare cause of acquired non-hormonal clitoromegaly.

Their occurrence usually follows a traumatic event, which is commonly represented by the female genital mutilation practice in case the cyst is located in the genital area. These cysts are extremely uncommon without prior history of genital surgery, infibulation, circumcision, or genital piercing [2].

The diagnosis of epidermoid cysts is usually done clinically and is based on the appearance of a soft, freely moveable cyst, often with a visible central punctum [1]. Radiologic imaging may be helpful to corroborate the clinical diagnosis; ultrasound evaluation shows these cysts as unilocular, hypoechoic, non-vascularized masses [3], and magnetic resonance imaging assessment could be useful, especially for determining the surgical excision strategy, given that such cysts have well-defined masses with high T2 signal background [3].

Since their degeneration into malignant squamous cell carcinoma is very rare, epidermal inclusion cysts generally do not require treatment but follow-up only [4]. They often stop growing after reaching a size of 1-5 cm in diameter [5]. However, if their growth causes severe discomfort and pain, a surgical approach should be considered as the first-line treatment [5].

While existing literature predominantly describes rare cases of spontaneous clitoral epidermal inclusion cysts, mostly in the paediatric population, this study presents a case of a large clitoral epidermal inclusion cyst in a post-menopausal woman without a history of genital trauma. Additionally, a review of all reported cases of spontaneous clitoral epidermal inclusion cysts in post-pubertal women was performed, contributing to the understanding of this rare condition.

CASE PRESENTATION

A 56-year-old woman referred to the gynaecological outpatient clinic for a routine check. She reported no history of genital trauma or mutilation, and had two vaginal deliveries. Apart from hypertension, she was in good health. The patient complained of a painless clitoral mass that had been progressively growing over the past two years, causing dyspareunia and psychological distress during sexual intercourses. There were no urinary complaints.

Pelvic examination revealed a mobile, soft, non-tender, well-defined clitoral mass measuring approximately 4cm, located above the urethra. The clitoral hood could not be retracted for visualization. No visible punctum or skin anomalies were noted (Fig. 1). Results of the general physical examination were normal and there were no other signs of hyperandrogenism.

An ultrasound scan characterized the mass as a 37x23mm non-vascularized anechoic cyst. Ultrasound examinations of the uterus, adnexal regions, and bladder were normal. Lab investigations were within normal limits. Discussions regarding the pros and cons of a wait and see approach took place, considering the patient's discomfort, and an elective surgery for the mass excision was organized with the patient's consent.

The mass was excised under spinal anaesthesia by both a gynaecologist and a plastic surgeon, ensuring a collaborative and comprehensive approach. They performed a longitudinal incision, and the cyst was removed with combination of sharp and blunt dissection, paying attention to avoiding the neurovascular bundle areas of the clitoris. The incision site was sutured with separate 3-0 absorbable sutures (Fig. 2), chosen for their appropriate dissolution rate and to minimize the need for suture removal in this sensitive area. The patient was discharged on the same day with anti-inflammatory medication and a wound-healing topic treatment, aimed at ensuring optimal recovery and minimizing discomfort. The post-operative period was completely uneventful.

The histopathology report revealed a 4cm cystic formation, with thin walls, lined with stratified keratinizing epithelium. These findings were instrumental in confirming the diagnosis of an epidermal inclusion cyst, guiding further management decisions.

A 6-week follow-up visit was scheduled, during which it was noted that the lesion was completely remodelled and adherent, without clinical signs of recurrence (fig. 3). Sensibility was preserved, and the patient referred a positive and satisfying return to her sexual life.

A signed informed consent for reporting and publishing this case was obtained from the patient.

DISCUSSION

When conducting a comprehensive search across major databases including PubMed, Medline, and Google Scholar, using key terms such as "clitoral", "clitoris", "epidermal cyst" and "epidermoid cyst", a total of 41 results were identified. Focusing on case reports and meta-analyses in English that involved adult and post-pubertal patients without history of genital trauma or surgery, 14 cases from 1999 until nowadays [3-14] were reviewed, in addition to the current case under analysis (Table 1). Among the reviewed cases, one occurred 18 months after caesarean-section delivery [6], and another during pregnancy [9]. In the latter case, the patient described a gradual and painless clitoral enlargement during pregnancy, with spontaneous regression within two months post-delivery. Intriguingly, five months later, the clitoral enlargement recurred and persisted during her second pregnancy.

Additionally, a case involving a 17-year-old patient highlighted a progressive regrowth of the cyst following vaginal delivery. Another unique case involved a 15-year-old patient experiencing a spontaneous recurrence seven years after iatrogenic drainage [8].

In the reviewed cases (Table 1), the mean age at the time of diagnosis was 27.5 ± 11.1 years old, with a mean time of growth of 21 ± 20.4 months. The mean cyst size was 4.8 ± 1.6 cm. Noteworthy findings include sexual-related symptoms reported by most adult patients, with dyspareunia observed in 25% and anorgasmia in 8.3%. Additionally, two patients presented urinary symptoms such as urinary frequency and voiding difficulty (16.6%), and a 47 years-old woman reported difficulties in walking due to the large cyst size (8.3%). Psychological distress was reported in 33% of the cases.

All patients underwent surgical excision using various techniques such as longitudinal incision, supra-meatal inverted-U-incision, and curvilinear incision, with no discernible clinical difference in post-operative recovery and patients' satisfaction. There has been only one case of post-operative

complication described – a secondary infection on the 2nd post-operative day, successfully treated with antibiotics

Follow-up was reported in 66,7% of cases (10/15 patients). Women attended follow-up visits from three months to four years after the surgery (median 9 months), with all reporting resolution of the symptoms. Anatomical restoration, preserved sexual function and maintained tactile sensation were consistently observed. Importantly, no cases of cyst recurrence were reported after surgery.

Conditions affecting the female external genitalia encompass vaginal (hymenial), para-urethral, and clitoral cysts. Among these, clitoral conditions include a spectrum of benign, malignant, and rarely metastatic lesions. Clitoromegaly is a rare condition that can be either congenital or acquired and can be attributable to hormonal or non-hormonal causes (Table 2).

The most common form of congenital clitoromegaly is associated with congenital adrenal hyperplasia (CAH), a condition that includes a group of genetic disorders affecting the adrenal glands and resulting in an increased production of adrenal-derived androgens. The excessive production of testosterone leads to various clinical signs and symptoms, including genitals' virilization, among which clitoromegaly is one of the most typical forms. More commonly, adult patients with ambiguous genitalia are found to have late-onset CAH, and present typical clinical signs such as hirsutism, irregular menses, or acne [15].

Acquired clitoromegaly is unusual and can be due to either hormonal or non-hormonal causes.

Hormonal causes of clitoromegaly often involve virilizing tumours, including Leydig-cell tumours, sex-cord stromal tumours, granulosa cell tumours or thecomas. Additional hormonal contributors encompass polycystic ovary syndrome and Cushing syndrome, and specific conditions observed during pregnancy, such as luteoma, hyperreaction luteinalis, and placental aromatase deficiency [9].

Non-hormonal causes, though rarer, encompass a spectrum of conditions. Non-tumoral forms include abscesses, diverticula, inguinal hernias, post-traumatic hematomas, or endometriomas. Tumoral forms range from benign entities like leiomyomas, pseudo-lymphomas, fibromas, angiokeratomas, hemangiomas, hemangiopericytomas, granular cell tumours, or neurofibromas, to malignancies such as endodermal sinus tumours, sarcomas, schwannomas, lymphomas, rhabdomyosarcomas, epithelioid hemangioendotheliomas or secondary metastasis (Table 3).

Acquired non-hormonal clitoromegaly is a rare phenomenon, and within this spectrum, the occurrence of epidermoid cyst in the clitoris is even more uncommon. Typically associated with genital mutilation in certain ethnic communities in Africa and West Asia, these cysts also manifest spontaneously without prior history of infibulation or piercing [16]. The evaluation of clitoromegaly poses a diagnostic challenge for clinicians due to the rarity and diverse aetiologies of this condition. Achieving an accurate differential diagnosis becomes imperative for guiding appropriate and effective treatment strategies.

CONCLUSIONS

While clitoral cysts are exceedingly rare, they play a crucial role in the differential diagnosis of clitoral hypertrophy. The evaluation of adult-onset clitoromegaly demands a comprehensive approach, involving meticulous physical examination, hormonal testing, and imaging studies. Surgical removal emerges as the preferred treatment for clitoral cyst due to the rarity of spontaneous regression and the potential recurrence after the procedure.

Studies have shown that timely diagnosis and treatment play a pivotal role in preventing negative outcomes, particularly in not-yet sexually active patients, while also contributing to the restoration of sexual well-being in those who have initiated sexual activity [17, 18].

Compliance with Ethical Standards

Authors contribution:

Federica Savasta contributed to the literature review and drafting of the manuscript.

Alessandro Libretti contributed to the conception of the case report and drafting of the manuscript.

Alberto Catinella was involved in patient care and revision of the article.

Antonio Taveggia was involved in patient care and revision of the article.

Libera Troia participated in the literature review and revising the article critically for important intellectual content.

Valentino Remorgida contributed to the conception of the case report and the revision of the article.

All authors approved the final submitted manuscript.

Funding: The authors declare that no funds, grants, or other support were received during the preparation of this manuscript..

Study registration: /.

Disclosure of Interests: The authors have no relevant financial or non-financial interests to disclose.

- Federica Savasta: none
- Alessandro Libretti: none
- Alberto Catinella: none
- Antonio Taveggia: none
- Libera Troia: none
- Valentino Remorgida: none.

Ethical Approval: /.

Informed consent: A signed informed consent for reporting and publishing this case was obtained from the patient.

Further Declarations:

The authors followed the EQUATOR Guidelines for reporting health research.

Simera I, Altman DG, Moher D, Schulz KF, Hoey J. Guidelines for reporting health research: the EQUATOR network's survey of guideline authors. PLoS Med. 2008 Jun 24;5(6):e139. doi: 10.1371/journal.pmed.0050139. PMID: 18578566; PMCID: PMC2443184.

Data sharing: Data regarding any of the subjects in the study has not been previously published.

References

1. Weir CB, St.Hilaire NJ. Epidermal Inclusion Cyst. 2022 Aug 8. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan–. PMID: 30335343.
2. Rouzi AA, Sindi O, Radhan B, et al. Epidermal clitoral inclusion cyst after type I female genital mutilation. *Am J Obstet Gynecol*. 2001 Sep;185(3):569-71. doi: 10.1067/mob.2001.117660. PMID: 11568779.
3. Johnson LT, Lara-Torre E, Murchison A, et. al. Large epidermal cyst of the clitoris: a novel diagnostic approach to assist in surgical removal. *J Pediatr Adolesc Gynecol*. 2013 Apr;26(2):e33-5. doi: 10.1016/j.jpag.2012.11.005. Epub 2013 Jan 20. PMID: 23337308.
4. Fux-Otta C, Fuster M, Ramos N, et al. Clitoromegaly due to an epidermal inclusion cyst: A case report. *Case Rep Womens Health*. 2022 Jul 18;35:e00432. doi: 10.1016/j.crwh.2022.e00432. PMID: 35898429; PMCID: PMC9309671.
5. Prasad I, Sinha S, Bharti S, et. al. Epidermal Inclusion Cyst of the Clitoris: A Case Report. *Cureus*. 2022 Sep 12;14(9):e29066. doi: 10.7759/cureus.29066. PMID: 36133501; PMCID: PMC9470212.
6. Oluwaferanmi E, Jha S. Clinical image-postpartum clitoral epidermoid cyst. *Int Urogynecol J*. 2022 Sep;33(9):2589-2590. doi: 10.1007/s00192-022-05177-7. Epub 2022 Apr 5. PMID: 35380211; PMCID: PMC9427885.
7. Mahmoudnejad N, Mohammadi Torbati P, Zadmehr A. Primary Epidermoid Cyst of the Clitoris in Adult Female Population: Three Case Reports and Introducing a Safe Surgical Approach. *Urol J*. 2020 Nov 4;18(3):343-346. doi: 10.22037/uj.v16i7.6348. PMID: 33159314.
8. DiCarlo-Meacham AM, Dengler KL, Snitchler AN, et al. Clitoral Epidermal Inclusion Cyst Leading to Anorgasmia: A Case Report and Literature Review. *J Pediatr Adolesc Gynecol*. 2020 Jun;33(3):321-323. doi: 10.1016/j.jpag.2020.01.150. Epub 2020 Feb 3. PMID: 32028052.
9. Hughes JW, Guess MK, Hittelman A et al. Clitoral epidermoid cyst presenting as pseudoclitoromegaly of pregnancy. *AJP Rep*. 2013 May;3(1):57-62. doi: 10.1055/s-0033-1334461. Epub 2013 Feb 7. PMID: 23943712; PMCID: PMC3699154.
10. Paulus YM, Wong AE, Chen B, et al. Preputial epidermoid cyst: an atypical case of acquired pseudoclitoromegaly. *J Low Genit Tract Dis*. 2010 Oct;14(4):382-6. doi: 10.1097/LGT.0b013e3181d85da9. PMID: 20885168.
11. Anderson-Mueller BE, Laudenschlager MD, Hansen KA. Epidermoid cyst of the clitoris: an unusual cause of clitoromegaly in a patient without history of previous female circumcision. *J Pediatr Adolesc Gynecol*. 2009 Oct;22(5):e130-2. doi: 10.1016/j.jpag.2008.10.006. Epub 2009 Jul 3. PMID: 19576807.
12. Fedele L, Fontana E, Bianchi S, et al. An unusual case of clitoromegaly. *Eur J Obstet Gynecol Reprod Biol*. 2008 Oct;140(2):287-8. doi: 10.1016/j.ejogrb.2008.02.008. Epub 2008 Apr 2. PMID: 18387729.
13. Linck D, Hayes MF. Clitoral cyst as a cause of ambiguous genitalia. *Obstet Gynecol*. 2002 May;99(5 Pt 2):963-6. doi: 10.1016/s0029-7844(02)01967-1. PMID: 11975977.
14. Schmidt A, Lang U, Kiess W. Epidermal cyst of the clitoris: a rare cause of clitorimegaly. *Eur J Obstet Gynecol Reprod Biol*. 1999 Dec;87(2):163-5. doi: 10.1016/s0301-2115(99)00096-2. PMID: 10597967.

15. Aaa El-Maouche D, Arlt W, Merke DP. Congenital adrenal hyperplasia. *Lancet*. 2017 Nov 11;390(10108):2194-2210. doi: 10.1016/S0140-6736(17)31431-9. Epub 2017 May 30. Erratum in: *Lancet*. 2017 Nov 11;390(10108):2142. PMID: 28576284.
16. Libretti A, Bianco G, Corsini C, et al. Female genital mutilation/cutting: going beyond urogynecologic complications and obstetric outcomes. *Arch Gynecol Obstet*. 2023 Oct;308(4):1067-1074. doi: 10.1007/s00404-023-06929-6. Epub 2023 Jan 27. PMID: 36703012.
17. Laganà AS, Vitale SG, Stojanovska L. et al., Preliminary results of a single-arm pilot study to assess the safety and efficacy of visnadine, prenylflavonoids and bovine colostrum in postmenopausal sexually active women affected by vulvovaginal atrophy. *Maturitas*. 2018 Mar;109:78-80. doi: 10.1016/j.maturitas.2017.12.015. Epub 2017 Dec 24. PMID: 29452786.
18. Vitale SG, Laganà AS, Nigro A et al., Peroxisome Proliferator-Activated Receptor Modulation during Metabolic Diseases and Cancers: Master and Minions. *PPAR Res*. 2016;2016:6517313. doi: 10.1155/2016/6517313. Epub 2016 Dec 28. PMID: 28115924; PMCID: PMC5225385.

Table 1 – Reported cases of adult-onset spontaneous epidermoid cyst of the clitoris

	Reference	Age (years)	Time of growth (months)	Size (cm)	Clinical presentation	Treatment	Follow-up (months)	
1	Current case	56	24	4	Dyspareunia and psychological distress	Surgical excision (longitudinal incision)	3	Well-being, no recurrence
2	Fux-Otta et al ⁴ , 2022	16	12	4	Emotional discomfort	Surgical excision	1 2	Sexual satisfaction, discomfort resolution, well-being
3	Prasad et al ⁵ , 2022	43	36	6	Dyspareunia	Surgical excision (curvilinear incision)	3	Asymptomatic, sexual discomfort resolution, no recurrence
4	Oluwaferanmi et al ⁶ , 2022	31, 18 months post partum	19	8	ND	Surgical excision	ND	Secondary infection, successfully treated with antibiotics
5	Mahmoudnejad et al ⁷ , 2021	45	6	5	Asymptomatic	Surgical excision (supra-meatal inverted-U-incision)	6	Sexual and tactile sensation preserved, no recurrence
		35	24	3	Dyspareunia and difficulties in voiding		1 2	Asymptomatic, no recurrence
		25	24	5	Urinary symptoms (urinary frequency and difficulties in voiding)		4 8	Asymptomatic, sexual well-being
6	Angela M. DiCarlo-Meacham et al ⁸ , 2020	15, recurred at 22	84	4	Anorgasmia	Surgical excision (transverse incision)	7	Orgasm regained

7	Hughes et al ⁹ , 2013	31, pregnant	5 post-partum	7	Psychological distress	Surgical excision	2	Asymptomatic, no recurrence
9	Johnson et al ³ , 2010	15	24	5	Asymptomatic, social anxiety	Surgical excision	3	Anatomy restoration, sexual and tactile sensation preserved, no recurrence
10	Paulus et al ¹⁰ , 2010	47	12	4	Pain	Surgical excision	ND	ND
11	Anderson-Mueller et al ¹¹ , 2009	17	Spontaneous discharge, recurrence	6	Pain	Surgical excision	ND	Asymptomatic
12	Fedele et al ¹² , 2007	22	10	7	Asymptomatic	Surgical excision (vertical incision)	ND	Anatomic restoration, preservation of sensibility
13	Linck et al ¹³ , 2002	28	ND	3	ND	Surgical excision	ND	ND
14	Schmidt et al ¹⁴ , 1999	16	8-10	2	ND	Surgical excision	18	No recurrence

Table 2 – Causes of clitoromegaly

Congenital	Acquired	
	Hormonal	Non hormonal
<ul style="list-style-type: none"> - CAH <ul style="list-style-type: none"> - Maternal hyperandrogenism: - Luteoma - Hyperreaction luteinalis - Virilizing tumors - Exogenous androgens intake 	<ul style="list-style-type: none"> - Polycystic ovary syndrome - Virilizing ovarian tumors <ul style="list-style-type: none"> - Leydig-cell tumor - Granulosa cell tumor - Thecomas - Sex-cord stromal tumors - Virilizing adrenal tumors <ul style="list-style-type: none"> - Adrenocortical carcinoma - Adrenocortical adenoma - Synthetic androgen exposure 	<ul style="list-style-type: none"> - Neurofibromatosis - Vulvar tumors, primary or metastatic - Hematomas - Inguinal hernias - Abscesses - Diverticula - Endometrioma

Table 3 – Vulvar tumors

Benign	Malign
Leiomioma	Endodermal sinus tumor
Pseudolymphoma	Sarcoma (liposarcoma)
Fibroma	Schwannoma
Angiokeratoma	Lymphoma
Hemangioma	Rhabdomyosarcoma
Hemangiopericytoma	Epithelioid hemangioendothelioma
Granular tumor cell	Metastasis
Neurofibroma	

Figure 1. – Clitoral cyst before surgery



Figure 2– Inspection of vulvar region following surgery



Figure 3 - Inspection of vulvar region at 6-week follow-up

