

The role of umbilical vein blood flow assessment in the prediction of foetal growth: a prospective observational cohort study

Daniele Farsetti^{1,2,*}, Moira Barbieri³, Francesca Pometti^{1,2}, Elena Magni^{4,5}, Barbara Vasapollo^{1,2}, Herbert Valensise^{1,2}, Tamara Stampalija^{4,5}

¹Department of Obstetrics and Gynecology, Policlinico Casilino, Rome, Italy.

²Department Surgical Sciences, University of Rome Tor Vergata, Rome, Italy.

³Clinica Mangiagalli, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, University of Milan, Milan, Italy.

⁴Department of Obstetrics and Gynecology, Unit of Fetal Medicine and Prenatal Diagnosis, Institute for Maternal and Child Health IRCCS Burlo Garofolo, Trieste, Italy.

⁵Department of Medicine, Surgery and Health Sciences, University of Trieste, Trieste, Italy.

DOI: 10.36129/jog.2024.S32

Objective. To evaluate the umbilical vein blood flow volume (UV-Q) in SGA fetuses and FGR, and to explore the correlation between UV-Q and foetal growth velocity (FGV). Secondly, the capacity of UV-Q and FGV in predicting adverse perinatal outcome (APO) and iatrogenic preterm birth were assessed.

Materials and Methods. 122 women were enrolled (64 SGA and 58 FGR according to Delphi consensus criteria). At the time of diagnosis foetal biometry and Doppler assessment, including absolute UV-Q and normalized for estimated foetal weight (UV-Q/EFW) and abdominal circumference (UV-Q/AC) were considered. The FGV was calculated from the difference between the EFW calculated in two consecutive sonographic evaluations. The pregnancies were followed until delivery and maternal-neonatal outcomes were collected.

Results. When compared to SGA and reference ranges, FGR

had significantly lower UV-Q, UV-Q/EFW, and UV-Q/AC. The FGV had a positive significant correlation with UV-Q ($r = 0.46$), UV-Q/AC ($r = 0.43$), and BW ($r = 0.56$).

The multivariable logistic regression analysis showed that UV-Q ≤ 0.65 MoM (aOR = 3.5) and FGV ≤ 0.63 MoM (aOR = 3.0) were independently associated with the occurrence of APO; UV-Q ≤ 0.60 MoM (aOR = 5.2) and FGV ≤ 0.63 MoM (aOR = 3.6) were independent predictors of iatrogenic preterm birth. This was true both for SGA and FGR.

Conclusions. The UV-Q might have a potential role in identifying fetuses with FGR and predicting foetal growth at the subsequent biometric evaluation. UV-Q and FGV are independent predictors of iatrogenic preterm birth and APO in a population of small fetuses, regardless of Delphi consensus criteria. These results encourage future studies on the predictive value of this parameter.