

SYSTEMATIC REVIEW AND META-ANALYSIS

Ablation and risk of recurrence in endometrioma: a systematic review and meta-analysis

Carlo **Ronsini** *, Irene **Iavarone**, Maria Giovanna **Vastarella**, Clorinda **Vitale**, Giada **Andreoli**, Marco **Torella**, Pasquale De **Franciscis**

Department of Woman, Child and General and Specialized Surgery, University of Campania "Luigi Vanvitelli", Naples, Italy

***Corresponding author:** Carlo **Ronsini**, M.D., Department of Woman, Child and General and Specialized Surgery, University of Campania "Luigi Vanvitelli", Naples, Italy.

Email: carlo.ronsini@unicampania.it

Doi: 10.36129/jog.2025.216

ABSTRACT

Objective. Ovarian endometriosis, or endometriomas, is a common manifestation and is typically treated via laparoscopic cystectomy. However, this method may compromise ovarian function. Ablation, involving thermal destruction of endometriosis cells, is emerging as an alternative. Despite some evidence suggesting less impact on ovarian function, there is a lack of robust data on the efficacy of ablation in preventing recurrence. This systematic review and meta-analysis aim to evaluate the recurrence rates of ablation compared to standard cystectomy for ovarian endometriomas.

Materials and Methods. This study followed PRISMA guidelines and was registered with PROSPERO (protocol number CRD549177). Comprehensive searches were conducted in PubMed, EMBASE, Scopus, Google Scholar, ClinicalTrials.gov, and the Cochrane Central Register up to May 2024. Inclusion criteria focused on studies involving patients with at least one ovarian endometrioma treated with ablation or cystectomy, reporting recurrence rates, and having a minimum follow-up of 12 months. The studies were assessed for quality using the Newcastle-Ottawa Scale. Data were analyzed using fixed-effect or random-effect models based on heterogeneity, with statistical significance set at $p < 0.05$.

Results. The search identified 58 articles, with 16 meeting the criteria for review. 5 studies, encompassing 395 patients, were included in the final analysis. 4 studies compared ablation and cystectomy. Recurrence rates varied, with ablation ranging from 0% to 37.7% and cystectomy from 0% to 22%. Meta-analysis revealed a non-significant trend toward higher recurrence rates with cystectomy (OR 1.99, 95% CI 0.95-4.16, $p=0.07$). The heterogeneity was low ($I^2=0\%$, $p=0.45$).

Conclusions. This systematic review and meta-analysis did not find a statistically significant difference in recurrence rates between ablation and cystectomy for treating ovarian endometriomas. However, there was a non-significant trend favoring ablation. Further randomized controlled trials are necessary to confirm these findings and to better understand the long-term efficacy and safety of ablation compared to cystectomy.

Key words

Endometriosis; ovarian endometrioma; ablation; laparoscopic cystectomy; recurrence rate.

INTRODUCTION

Endometriosis, a condition affecting about 10% of women of childbearing age, involves the displacement of endometrial tissue outside the uterus [1-2]. Ovarian endometriosis, also known as endometriomas, is its most frequent presentation and involves several therapeutic approaches. While laparoscopic cystectomy, also called stripping, is the current standard, surgery can cause damage to ovarian tissue, diminishing its endocrinological and reproductive potential [3]. Various approaches have been studied to minimize this risk [4]. Among these, ablation, which involves destroying the endometriosis cells by applying energy from different sources, but which has in common the thermal damage done to the endometrioma, has found increasing use in recent years [5]. Although it is now considered an alternative method to laparoscopic stripping, the scarcity of prospective studies raises questions about its efficacy in terms of risk of disease recurrence. Recently, a meta-analysis has shown its minor impact on ovarian function [6].

In contrast, solid data on its efficacy over time in controlling the development of new endometriomas are lacking in the literature. Moreover, the lack of standardization of the technique may make it even more challenging to understand its efficacy fully. This is why we wanted to collect all the data to date in the literature on this topic. This systematic review and meta-analysis aim to assess ablation's recurrence outcomes compared to standard cystectomy for ovarian endometriomas.

MATERIAL AND METHODS

The methods for this study were specified a priori based on the recommendations in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement [9]. We registered the Review for meta-analysis on the PROSPERO site with protocol number CRD549177.

Search Method

We performed systematic research for records about the use of sclerotherapy in managing ovarian endometriomas in PubMed, EMBASE, Scopus, Google Scholar, Clinical-trials.gov, and the Cochrane Central Register of Controlled Trials in May 2024. We did not restrict country or year of publication and considered only entirely English-published studies. We adopted the following string of idioms in each database to identify studies fitting to our review's topic: "Endometriosis and Ablation".

Study Selection

Study selection was made independently by G.A. and M.G.V. In case of discrepancy, C.R. decided on inclusion or exclusion. Inclusion criteria were: (1) studies that included patients with at least one ovarian endometrioma, treated with Ablation and/or cystectomy; (2) studies reporting the outcome of interest: Recurrence Rate (RR); (3) Studies with at least 12 months of follow-up; (4) peer-reviewed articles, published originally. We excluded non-original studies, pre-clinical trials, animal trials, abstract-only publications, and articles in languages other than English. If possible, the authors of studies that were published as conference abstracts were tried to be contacted via e-mail and asked to provide their data. We mentioned the studies selected and all reasons for exclusion in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flowchart (Fig. 1). We assessed all included studies regarding potential conflicts of interest.

Statistical analysis

Heterogeneity among the studies was tested using the Chi-square test and I-square tests [7]. The Odds Ratio (OR) and 95% confidence intervals (CI) were used for dichotomous variables. Fixed-effect models conducted statistical analysis without significant heterogeneity ($I^2 < 50\%$), or random-effect models if $I^2 > 50\%$. Recurrence rate (RR) was used as clinical outcomes. In each study, RR was defined as the percentage of recurrence till the last follow-up. Chi-square tests were used to compare continuous variables. Review Manager version 5.4.1 (REVman 5.4.1) and IBM Statistical Package for Social Science (IBM SPSS vers 25.0) for MAC were used for statistic calculation. For all performed analyses, a p-value < 0.05 was considered significant.

Quality assessment

We assessed the quality of the included studies using the Newcastle–Ottawa scale (NOS) [8]. This assessment scale uses three broad factors (selection, comparability, and exposure), with the scores ranging from 0 (lowest quality) to 8 (best quality). Two authors (CR and II) independently rated the study's quality. Any disagreement was subsequently resolved by discussion or consultation with PDF. We reported NOS Scale in Appendix.

We used a funnel plot analysis to assess publication bias. We used Egger's regression test to determine the asymmetry of funnel plots (Appendix).

Risk of Bias

The RCTs and prospective cohort studies were separately assessed, and the risk of bias in these studies was low or moderate. Saito et al. [11] only included nulliparae in the ablation group and reported bigger dimensions of endometrioma in the ablation group. In addition, he has the shortest follow-up and has not reported any recurrence events in either group during these twelve months. Haghgoo et al. included 30 patients with unilateral endometrioma and 30 with bilateral endometrioma [9]. Candiani et al. used a CO2 fiber laser for cyst vaporization [12], whereas argon is employed commonly.

RESULTS

Studies' Characteristics

After the database search, 58 articles matched the search criteria. After removing records with no full text, duplicates, and wrong study designs (e.g., reviews), 16 were eligible. 5 matched the inclusion criteria and were included in the systematic review. 4 were comparative studies between the Ablation technique and laparoscopic stripping and were included in quantitative analysis (Fig 1). The countries where the studies were conducted, the publication year range, the studies' design, Follow-up months, and the number of participants are summarized in Table 1.

NOS [8] (Appendix) assessed the quality of all studies. Overall, the publication years ranged from 2011 to 2021. In total, 395 patients with endometrioma were enrolled: 215 were treated with ablation and 180 with laparoscopic stripping.

Outcomes

The review included 395 patients. All 5 selected studies presented RR data. Overall, the RR ranged from 0 to 37.7% in the ablation group and from 0 to 22.0% in the stripping group. The follow-up period ranged from 12 to 64 months on average. Those results are summarized in Table 2.

By alphabetic, Candiani et al [12] reported a RR of 6.3 vs 4.9 for ablation compared to stripping in 29 months of Follow-up ($p=0.74$). Carmona et al [10] reported the oldest series with the highest RR in both arms (22% vs 37%, $p=0.4$) and longest Follow-Up (64 months). On the contrary, Chen et al

[13] is the newest one, with RR 4.4% vs 16.7% ($p=0.11$) and 31 months of observation. Haghgoo et al [9] reported the only single-arm trial with no recurrence reported after 15 months. Finally Saito et al [11] did not observed recurrence in both arms.

Ablation Procedure

Haghgoo et al. avoided hot energy devices, as cautery, on ovaries for ablation [9]. In Carmona et al. study, the vaporization of the cyst's internal wall was performed through CO₂ laser at 30 W/cm² power density [10]. Saito et al. performed vaporization using bipolar current forceps (35 W) on the internal wall [11]. Candiani et al. used a CO₂ fiber laser in a "one-step" procedure [12]. In Chen et al. study, bipolar forceps were applied on the internal wall at 30W until the color of the cyst turned white [13]. The average duration of contact between the forceps and the lesion was approximately 1 second [13]. In all cases, a biopsy was performed before proceeding with vaporization [9-13].

Meta-analysis

The 4 studies comparing ablation and stripping were enrolled in the meta-analysis. A total of 337 patients were analyzed. 157 patients in the ablation arm were compared with 180 patients who underwent cystectomy, exploring RR outcome. 23 recurrences occurred in the ablation group and 14 in the stripping. Because of low heterogeneity ($I^2=0\%$; $p = 0.45$), the fixed-effects model was applied. The Cystectomy group showed a slightly non-significant higher risk for recurrence than the ablation arm (RR 1.99 [95% CI 0.95-4.16] $p=0.07$). Fig 2.

DISCUSSION

Main Findings

This systematic review and meta-analysis included five studies, four cohort studies, and one randomized trial. The qualitative data analysis could not show a statistically significant difference between the ablation technique and laparoscopic stripping based on recurrence ($p=0.07$). However, stripping seems to show a worsening trend with an OR per recurrence of 1.99 and a 95% CI slightly including neutrality (0.95-4.16). This could be confirmed as the sample size increases. Moreover, the largest weighting (51.1%) is represented by Carmona et al [10] which has the advantage of being the only randomised clinical trial and the disadvantage of being the oldest included study, risking being flawed by a technological backwardness, which may undermine the efficacy of the ablative technique.

Comparison with existing literature.

Since its first publication, ablation has been an attractive alternative for treating endometriomas. This is because the relationship between endometriosis and infertility is an intrinsic one, worsened by all treatment episodes of a surgical nature. Cyst removal inevitably results in a reduction in the patient's reproductive and endocrinological potential. Much scientific society has struggled to minimize this impact, on the one hand, by attempting to optimize post-stripping coagulation techniques [3] and, on the other hand in, seeking alternative approaches for treating endometrioma, such as sclerotherapy [4]. In the same vein, all alternatives provided to stripping, including ablation, have always been weighed first in assessing their impact on fertility. Recently, Zhang et al [5] published a meta-analysis on pre- and postoperative differences in AMH and Antral Follicle Count (AFC) determined by ablation and stripping. The 294 patients enrolled in this meta-analysis showed a lower AFC in the stripping arm, both in the immediate postoperative period (mean differences [MD], 1.33; 95% credible interval, 2.15 to 0.51; I^2 1/4 57%), and at 6-month follow-up (MD, 1.93; 95% credible interval, 2.40 to 1.45; I^2 1/4 0%). The intragroup comparisons of AMH levels supported negative effects on ovarian reserve of both cystectomy (MD, 1.26; 95% credible interval, 1.64 to 0.88; I^2 1/4 45%) and ablation (MD, 0.70; 95% credible interval, 1.07 to

0.32; I2 1/4 0%). These data support the evidence from another systematic review on the use of cystectomy [14] and a meta-analysis [15] conducted by two independent groups. A group in our analysis also published data from the same series on this subject, with a more significant impact of cystectomy on AMH and AFC [16]. On the other hand, much rarer are the papers that focus on ablation's efficacy regarding recurrence risk. In this scenario, to our knowledge, our meta-analysis stands as the only meta-analysis focusing on this topic.

Clinical Implication

Once the lower impact on patients' fertility has been established, it is also essential to weigh up the effectiveness of the ablative technique over time. This is why, in our opinion, the evidence derived from our work supports the therapeutic choices for treating patients with endometriosis.

Endometriosis is a chronic condition where gaining time between treatment episodes is crucial to optimizing treatment. To date, surgery remains the diagnostic gold standard, but this contrasts with the need to be minimally invasive. This discrepancy often creates diagnostic delays that condition the severity of the clinical presentation [17]. On the one hand, we need to optimize diagnosis by identifying biomarkers, such as liquid biopsy [18], that can intercept our patients before organic progression, and on the other hand, we need to improve our therapeutic options to chronicle the disease with as little morbidity as possible. Fortunately, the pharmacological landscape has recently expanded with new drugs such as Relugolix, which have shown promising results in controlling symptoms [19]. Surgery should go hand in hand with adapting to the growing number of therapeutic options. In this scenario, we believe standardization of the ablative technique would also be fundamental to improve its reproducibility.

Strengths and Limitations

Our study is the first meta-analysis to evaluate whether cystectomy or ablation results in higher endometrioma RR. Unfortunately, very few studies on this subject limit our case series to 395. Our results were not statistically significant, even though they showed a clear trend against cystectomy. Another limitation is the lack of standardization in ablative techniques and hemostatic approaches in the case of stripping. This attempt to compare the two methods is more ambitious.

Furthermore, studies that have used bipolar coagulation have not reported any data on its intensity. A further limitation is the possibility of bilateral neoformations, which were included in the studies as a single patient, even though it is assumed that the double procedure exposes one to a double risk of recurrence. Finally, no data have been reported regarding spillage during treatment, which may promote intraabdominal spillage of endometriosis tissue and increase the chances of recurrence [20-21].

Finally, a final point should be made that our review does not take into account additional outcomes that may differentiate the two techniques, such as the risk of postoperative pain [22] or the risk of malignant transformation [23].

CONCLUSIONS

Our study failed to show a statistically significant ($p=0.07$) increased safety profile of ablation compared to cystectomy in terms of RR. However, the data show a clear trend with almost doubled risk of recurrence in patients undergoing laparoscopic cystectomy (OR 1.99 CI 95% 0.95-4.16). Further randomised trials may support or refute this trend.

Author contributions

Carlo Ronsini: Conceptualization; Formal analysis; Methodology; Project administration; Software; Supervision; Validation; Writing - original draft; Writing - review & editing, Irene Iavarone: Data curation; Writing - review & editing, Maria Giovanna Vastarella: Data curation; Investigation; Project administration; Software; Supervision, Clorinda Vitale: Data curation, Investigation, Giada Andreoli:

Data curation; Formal analysis; Marco Torella: Supervision; Validation; Visualization; Pasquale de Franciscis: Supervision; Validation; Visualization.

Funding

None.

Study registration

We registered the Review for meta-analysis on the PROSPERO site with protocol number CRD549177.

Competing interests statement

The authors of the manuscript have nothing to disclosure about it. ETHICAL APPROVAL Not applicable.

Informed consent

Not applicable.

Data sharing

No dataset was made to collect data for this article. Literature data are present in the reference list.

REFERENCES

REFERENCES

1. Burney RO, Giudice LC. Pathogenesis and pathophysiology of endometriosis. *Fertil Steril*. 2012 Sep;98(3):511-9. doi: 10.1016/j.fertnstert.2012.06.029. Epub 2012. PMID: 22819144; PMCID: PMC3836682.
2. Sanchez AM, Viganò P, Somigliana E, Panina-Bordignon P, Vercellini P, Candiani M. The distinguishing cellular and molecular features of the endometriotic ovarian cyst: from pathophysiology to the potential endometrioma-mediated damage to the ovary. *Hum Reprod Update*. 2014;20(2):217-30. doi: 10.1093/humupd/dmt053. Epub 2013 Oct 14. PMID: 24129684.
3. Riemma G, De Franciscis P, La Verde M, Ravo M, Fumiento P, Fasulo DD, Della Corte L, Ronsini C, Torella M, Cobellis L. Impact of the hemostatic approach after laparoscopic endometrioma excision on ovarian reserve: Systematic review and network meta-analysis of randomized controlled trials. *Int J Gynaecol Obstet*.;162(1):222-232. doi: 10.1002/ijgo.14621. Epub 2022 Dec 26. PMID: 36503998.
4. Ronsini C, Iavarone I, Braca E, Vastarella MG, De Franciscis P, Torella M. The Efficiency of Sclerotherapy for the Management of Endometrioma: A Systematic Review and Meta-Analysis of Clinical and Fertility Outcomes. *Medicina (Kaunas)*. 2023;59(9):1643. doi: 10.3390/medicina59091643. PMID: 37763762; PMCID: PMC10535205.
5. Jee BC. Efficacy of ablation and sclerotherapy for the management of ovarian endometrioma: A narrative review. *Clin Exp Reprod Med*.;49(2):76-86. doi: 10.5653/cerm.2021.05183. Epub 2022 May 4. PMID: 35698769; PMCID: PMC9184881.
6. Zhang Y, Zhang S, Zhao Z, Wang C, Xu S, Wang F. Impact of cystectomy versus ablation for endometrioma on ovarian reserve: a systematic review and meta-analysis. *Fertil Steril*. 2022 Dec;118(6):1172-1182. doi: 10.1016/j.fertnstert.2022.08.860. Epub 2022. PMID: 36334993.

7. Chaimani A, Higgins JP, Mavridis D, Spyridonos P, Salanti G. Graphical tools for network meta-analysis in STATA. *PLoS One*. 2013;8(10):e76654. doi: 10.1371/journal.pone.0076654. PMID: 24098547; PMCID: PMC3789683.
8. Kansagara, D.; O'Neil, M.; Nugent, S.; Freeman, M.; Low, A.; Kondo, K.; Elven, C.; Zakher, B.; Motu'apuaka, M.; Paynter, R.; et al. Benefits and Harms of Cannabis in Chronic Pain or Post-traumatic Stress Disorder: A Systematic Review [Internet]. Washington (DC): Department of Veterans Affairs (US); 2017. [Table], Quality Assessment Criteria for Observational Studies, Based on the Newcastle-Ottawa Scale. Available online: <https://www.ncbi.nlm.nih.gov/books/NBK476448/table/appc.t4/> (accessed on 12 August 2017).
9. Haghgoo A, Shervin A, Chaichian S, Ghahremani M, Mehdizadeh Kashi A, Akhbari F. Increasing trend of serum antimullerian hormone level after long term follow up of endometrioma resection. *Journal of Endometriosis and Pelvic Pain Disorders*. 2021;13(2):98-103. doi:10.1177/2284026521990465
10. Carmona F, Martínez-Zamora MA, Rabanal A, Martínez-Román S, Balasch J. Ovarian cystectomy versus laser vaporization in the treatment of ovarian endometriomas: a randomized clinical trial with a five-year follow-up. *Fertil Steril*. 2011 Jul;96(1):251-4. doi: 10.1016/j.fertnstert.2011.04.068. Epub 2011. PMID: 21575941.
11. Saito N, Yamashita Y, Okuda K, Kokunai K, Terai Y, Ohmichi M. Comparison of the impact of laparoscopic endometriotic cystectomy and vaporization on postoperative serum anti-Mullerian hormone levels. *Asian J Endosc Surg*. 2018 Feb;11(1):23-29. doi: 10.1111/ases.12412. Epub 2017. PMID: 28786171.
12. Candiani M, Ottolina J, Schimberni M, Tandoi I, Bartiromo L, Ferrari S. Recurrence Rate after "One-Step" CO2 Fiber Laser Vaporization versus Cystectomy for Ovarian Endometrioma: A 3-Year Follow-up Study. *J Minim Invasive Gynecol*. 2020 May-Jun;27(4):901-908. doi: 10.1016/j.jmig.2019.07.027. Epub 2019. PMID: 31377455.
13. Chen J, Huang D, Zhang J, Shi L, Li J, Zhang S. The effect of laparoscopic excisional and ablative surgery on ovarian reserve in patients with endometriomas: A retrospective study. *Medicine (Baltimore)*. 2021;100(7):e24362. doi: 10.1097/MD.00000000000024362. PMID: 33607770; PMCID: PMC7899828.
14. Raffi F, Metwally M, Amer S. The impact of excision of ovarian endometrioma on ovarian reserve: a systematic review and meta-analysis. *J Clin Endocrinol Metab*.;97(9):3146-54. doi: 10.1210/jc.2012-1558. Epub 2012 Jun 20. PMID: 22723324.
15. Candiani M, Ferrari SM, Salmeri N, Dolci C, Villanacci R, Bartiromo L, Schimberni M, Tandoi I, Papaleo E, Ottolina J. CO2 fiber laser vaporization for endometrioma treatment results in preserved ovarian responsiveness and improved embryo quality in infertile women undergoing ART. *Minerva Obstet Gynecol*. 2023;75(4):348-356. doi: 10.23736/S2724-606X.22.05188-0. Epub 2022 Oct 18. PMID: 36255166.
16. Agarwal SK, Chapron C, Giudice LC, Laufer MR, Leyland N, Missmer SA, Singh SS, Taylor HS. Clinical diagnosis of endometriosis: a call to action. *Am J Obstet Gynecol*. 2019;220(4):354.e1-354.e12. doi: 10.1016/j.ajog.2018.12.039. Epub 2019 Jan 6. PMID: 30625295.
17. Ronsini C, Fumiento P, Iavarone I, Greco PF, Cobellis L, De Franciscis P. Liquid Biopsy in Endometriosis: A Systematic Review. *Int J Mol Sci*. 2023;24(7):6116. doi: 10.3390/ijms24076116. PMID: 37047088; PMCID: PMC10094565.

18. Giudice LC, As-Sanie S, Arjona Ferreira JC, Becker CM, Abrao MS, Lessey BA, Brown E, Dynowski K, Wilk K, Li Y, Mathur V, Warsi QA, Wagman RB, Johnson NP. Once daily oral relugolix combination therapy versus placebo in patients with endometriosis-associated pain: two replicate phase 3, randomised, double-blind, studies (SPIRIT 1 and 2). *Lancet*. 2022;399(10343):2267-2279. doi: 10.1016/S0140-6736(22)00622-5. Erratum in: *Lancet*. 2022 Aug 27;400(10353):660. PMID: 35717987.
19. Chen Y, Liu X, Guo SW. Preoperative and perioperative intervention reduces the risk of recurrence of endometriosis in mice caused by either incomplete excision or spillage and dissemination. *Reprod Biomed Online*. 2021;43(3):379-393. doi: 10.1016/j.rbmo.2021.04.017. Epub 2021 Apr 29. PMID: 34330642.
20. Iavarone I, Greco PF, La Verde M, Morlando M, Torella M, de Franciscis P, Ronsini C. Correlations between Gut Microbial Composition, Pathophysiological and Surgical Aspects in Endometriosis: A Review of the Literature. *Medicina (Kaunas)*. 2023;59(2):347. doi: 10.3390/medicina59020347. PMID: 36837548; PMCID: PMC9962646.
21. Ronsini C, Mosca L, Iavarone I, Nicoletti R, Vinci D, Carotenuto RM, et al. Oncological outcomes in fertility-sparing treatment in stage IA-G2 endometrial cancer. *Front Oncol*. 2022 Sep 16;12:965029. doi: 10.3389/fonc.2022.965029.
22. Buzzaccarini G, Török P, Vitagliano A, Petousis S, Noventa M, Hortu I, et al. Predictors of Pain Development after Laparoscopic Adnexectomy: A Still Open Challenge. *J Invest Surg*. 2022 Jun;35(6):1392-1393. doi: 10.1080/08941939.2022.2056274
23. Ferrari F, Giannini A. Approaches to prevention of gynecological malignancies. *BMC Womens Health*. 2024 Apr 23;24(1):254. doi: 10.1186/s12905-024-03100-4.

Table 1. Characteristics of included studies.

Author, year of publication	Country	Period of enrollmen t	Study design	No. of participant s	Ablation	Cystecto my
Haghighi et al. 2021 [9]	Iran	2017-2019	Prospective Monocenter Cohort	58	0	58
Carmona et al. 2011 [10]	Spain	N/A	Prospective Monocenter Randomize d	74	38	36
Saito et al. 2017 [11]	Japan	2011-2013	Prospective Monocenter Cohort	62	28	34
Candiani et al. 2019 [12]	Italy	2015-2018	Prospective Monocenter Cohort	125	61	64
Chen et al. 2021 [13]	China	2016	Retrospectiv e Monocenter Cohort	76	30	46

FU: follow-up.

Table 2. Outcomes.

Author, year of publication	Cystectomy RR (%)	Ablation RR (%)	P-value	Median FU period (months)
-----------------------------	-------------------	-----------------	---------	---------------------------

Single-arm studies

Haghighi et al. 2021 [9]	-	0.0	N/A	15.0
--------------------------	---	-----	-----	------

Comparative studies

Carmona et al. 2011 [10]	22.0	37.0	0.4	64.0
--------------------------	------	------	-----	------

Saito et al. 2017 [11]	0.0	0.0	N/A	12.0
------------------------	-----	-----	-----	------

Candiani et al. 2019 [12]	6.3	4.9	0.74	29.0
---------------------------	-----	-----	------	------

Chen et al. 2021 [13]	4.4	16.7	0.11	31.38
-----------------------	-----	------	------	-------

RR: recurrence rate.

Figure 1. Flowchart.





Fig 2. Forrest Plot



APPENDIX

Appendix A. Newcastle-Ottawa Scale.

Author, year of publication	Country	Selection	Comparability	Exposure	Total
-----------------------------------	---------	-----------	---------------	----------	-------

Single-arm studies

Haghighi et al. 2021	Iran	3	1	1	5
----------------------	------	---	---	---	---

Comparative studies – Ablation vs. Cystectomy

Carmona et al. 2011	Italy	3	2	2	7
Saito et al. 2017	Japan	2	1	2	5
Candiani et al. 2019	Italy	2	1	3	6
Chen et al. 2021	China	2	2	3	7

Appendix-B Funnel Plot

