



Italian Journal of Gynæcology & Obstetrics

June 2026 - Vol. 38 - N. 2 - Quarterly - ISSN 2385 – 0868

Antioxidant therapy in endometriosis treatment: systematic review

Laura Pivazyan¹, Eva Nakhapetyan², Sapiyat Isaeva¹, Veronika Tarlakyán^{3,*}, Anastasia Laevskaya⁴, Eduard Ayryan³, Valeriia Seregina¹

¹FSBI “National medical Research Center for Obstetrics, Gynecology and Perinatology Named After Academician V.I.Kulakov” Ministry of Health of the Russian Federation, Moscow, Russia.

²Pirogov Russian National Research Medical University, Moscow, Russia.

³I.M. Sechenov First Moscow State Medical University (Sechenov University), Moscow, Russia.

⁴Botkin Hospital, Moscow, Russia.

ARTICLE INFO

History

Received: 14 April 2025

Received in revised form: 11 June 2025

Accepted: 17 July 2025

Available online: 22 June 2026

DOI: 10.36129/jog.2025.239

Key words

Endometriosis; antioxidants; treatment; oxidative stress.

*Corresponding author: Veronika Tarlakyán,

MBBS. I.M. Sechenov First Moscow State Medical University (Sechenov University),

8-2 Trubetskaya Street,

Moscow, Russian Federation 119991.

Email: vtarlakyan00@mail.ru.

ORCID: 0000-0003-1163-2874.

ABSTRACT

Objective. To summarize current knowledge on the effectiveness of antioxidant therapy in endometriosis treatment.

Materials and Methods. A systematic review was conducted per PRISMA guidelines and registered in PROSPERO 2023 CRD42023454705. Studies published until November 2024 were identified through PubMed, The Cochrane Library, ClinicalTrials.gov, Google Scholar, and MEDLINE. COVIDENCE software was used for screening. Risk of bias was assessed using the Cochrane Handbook.

Results. Out of 512 studies, 11 were included in the systematic review. Endometriotic cysts weight and volume are dose-dependent parameters that are significantly lower in antioxidant treatment groups ($p < 0.05$). Neither histological cell scores nor trichrome fibrosis scores showed statistically significant differences among treatment and control groups ($p > 0.05$) in 2 out of 3 studies. Significantly lower levels of total oxidant status (TOS), nitric oxide (NO) and oxidative stress index (OSI) are evaluated in the antioxidant group compared to control. However, no significant differences were observed in malondialdehyde (MDA), superoxide dismutase (SOD) and catalase (CAT) levels. The number of follicles was significantly increased, and the atretic follicles number was significantly decreased after therapy ($p < 0.05$). The IVF, cleavage, blastocyst formation rates and blastocyst number were significantly higher in treatment group compared the control.

Conclusions. Antioxidants may be considered as a possible component of endometriosis therapy to potentially enhance fertility outcomes and slow disease progression, though current evidence is preliminary and requires further validation.

INTRODUCTION

Endometriosis is an estrogen-dependent inflammatory gynaecological disease defined by the

presence of endometrial-like mucosa outside the uterine cavity. The pathogenesis of endometriosis is supported by several theories, such as immunological, implantation (Sampson’s theory), dy-

sontogenetic, dissemination, metaplastic, genetic, hormonal, *etc.* [1]. Endometriosis has an impact on fertility, affecting the ovarian reserve, embryo quality, implantation and normal anatomical structure of the reproductive organs and surrounding tissues. However, the mechanisms leading to endometriosis-associated infertility are not fully understood. Currently, the role of oxidative stress (OS) leading to iron metabolism disorders in the pathogenesis of endometriosis is widely discussed. Endometriotic lesions are resistant to ferroptosis – iron-mediated programmed non-apoptotic cell death –, which allows their implantation in the peritoneal cavity [2]. There is a need for further research in this area due to its high relevance, theoretical and practical importance. A microenvironment with a high level of reactive oxygen species (ROS), free radicals and iron is created as a result of cyclic changes in ectopic endometriotic lesions, which increases their adhesion and the progression of the disease.

The presence of ROS in cells is a physiological process due to their formation in normal oxidative metabolism. They control the ovarian cycle, steroidogenesis and ovulation [3]. However, the imbalance between free radicals and the antioxidant system causes oxidative stress, leading to a reduction in oocyte quality [4].

In this systematic review, we observe the efficiency of antioxidants supplementation in endometriosis treatment.

MATERIALS AND METHODS

Study design

Our systematic review was conducted and reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement [5].

The present systematic review has been registered in the PROSPERO international prospective registry of systematic reviews by the National Institute of Health Research (NIHR). The registration number is PROSPERO 2023 CRD42023454705.

Search strategy

To identify relevant articles, we conducted an electronic database search using several databases: PubMed, Google Scholar, ClinicalTrials.gov, Cochrane library to identify studies using key words and MeSH terms. The date of the last screening was November 26, 2024. Using the advanced search tool

on PubMed, the following combination of key words was used: ((endometriosis) OR (endometrioma)) AND (antioxidants) AND (oxidative stress) AND (treatment). No filters or limits were used. Additionally, the search was conducted using MeSH terms (endometriosis [MeSH Terms]) AND ((antioxidants [MeSH Terms]) AND (treatment [MeSH Terms]) AND (oxidative stress [MeSH Terms])).

The Cochrane Library electronic database search strategy was conducted. The combination of the search was as follows: ((endometriosis) OR (endometrioma)) AND (antioxidants) AND (oxidative stress) AND (treatment). No filters or limits were used. MeSH terms were also screened (MeSH descriptor: [Endometriosis] explode all trees and with qualifier(s): [antioxidants - MeSH]).

The search was also conducted in the ClinicalTrials.gov electronic database using an advanced search combination: endometriosis | antioxidants.

Study selection

For search conducting and further screening COVidence software was used. To ensure the quality and accuracy of the search results, two investigators performed the search independently. After the initial search, all articles were reviewed based on their titles and abstracts. The full texts of the studies that appeared to be appropriate according to their titles and abstracts were reviewed. Potential trials were also identified by searching the reference lists of the eligible trials. We included randomized (RCTs) and non-randomized clinical trials. Only articles written in English were included. Abstracts from congresses and unpublished articles were not included. As this is a review of published studies, Institutional Review Board (IRB) approval was not sought.

Two investigators (E.N., I.S) independently read the full texts of the preselected articles to verify their eligibility. Any studies with duplicate records were excluded. To minimize potential bias during the review process, any disagreements about the inclusion or exclusion of preselected studies were resolved with the help of a third author (A.L).

Inclusion criteria

The inclusion criteria specified autograft endometriosis mice or rat models and women with endometriosis related infertility, receiving antioxidant therapy.

Studies that described high levels of oxidative stress markers due to non endometriosis-related




Figure 1. PRISMA flow chart diagram.

reasons, phytoalexins, antioxidant decoctions as a therapy were excluded.

Data extraction and quality assessment

The studies included were independently collected by two authors (E.N., I.S) using a standardized data extraction procedure. We obtained the following characteristics from our studies: study design, type of animal model, types of antioxidants and regimens used, and the number of patients in each group and the follow-up duration.

The analysis in animal models was aimed to evaluate the level of oxidative stress markers, embryo and oocyte quality, implant weight, volume and histological cell scores of endometriotic lesions after antioxidant therapy.

The analysis of human studies was aimed to establish the pregnancy outcomes in addition to previously mentioned parameters.

Risk of bias was assessed for each included study using the Cochrane Handbook for Systematic Reviews of Interventions [6]. Two reviewers (V.T., A.L) independently assessed the quality of the selected studies. A third investigator (L.P) was involved in

the case of inconsistencies. In accordance with the Cochrane Handbook for Systematic Reviews of Interventions, the RoB 2 tool [7] was used to assess the risk of bias for randomized controlled trials and ROBINS-I [8] for non-randomized trials, SYRCLE's RoB tool for animal model studies [9].

RESULTS

Summary of included studies

The study selection process is illustrated by the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) flow chart diagram (Figure 1).

A total of 512 publications were identified through an electronic database search of PubMed, Google Scholar, and ClinicalTrials.gov., Cochrane library. Of these, 425 studies were screened for title and abstract. 388 were excluded, 38 were selected for eligibility assessment. After screening the full-text articles, 27 were excluded for failing to meet the eligibility criteria. The references of the selected studies were additionally searched for other eligible studies and 7 studies were identified. Finally, 11 trials that met the criteria were included in the systematic review. A total of 73 rats, 14 mice and 759 women from ten trials were analysed. The detailed summary of the studies analysis is shown in Tables 1, 2.

Antioxidants effectiveness in animal population

In 4 studies, endometriosis was induced by transplanting autologous uterine tissue onto the peritoneal wall of female Wistar albino rats [10, 11] and female NMRI mice [12, 13].

The vitamin C effectiveness in animal population

Erten *et al.* [10] and Durak *et al.* [11] evaluated the efficacy of vitamin C in the prevention and regression of endometriotic implant development in an experimentally induced autografted endometriosis rat model. Implant weight and volume, histological cell scores and trichrome fibrosis scores were measured. Both authors suggest that histological scores are independent of vitamin C dosage, in contrast to implant volume. However, the exact reason for their insensitivity is not mentioned. It could either be due to the small number of animals included or the lack of direct effects.

Hoorsan *et al.* [12] conducted a study on the efficacy of vitamin C endometriosis treatment in the NMRI mouse model. In contrast to the previous two stu-

Table 1. Description of selected studies included in the review (rats and mice).

First author, year of publication	Title	Population	Follow up period	Intervention	Comparison	Outcomes
Ozlem Ulas Erten et al., 2016 [10]	Vitamin C is effective for the prevention and regression of endometriotic implants in an experimentally induced rat model of endometriosis	Female Wistar Albino Rats (n = 33) Weight = 209g-270g A group (n = 11) B group (n = 11) C group (n = 11)	42 days	Surgical induced endometriosis (autograft model) A group: 1 st operation + intravenous vitamin C 500mg/kg every 2 days B group: 1 st operation; 2 nd operation + intravenous vitamin C 500 mg/kg every 2 days C group: 1 st operation; 2 nd operation	Vitamin C vs no vitamin C in all groups	Implant volume at the 2 nd operation (mm ³) Implant volume at the 3 rd operation (mm ³) Weight 1 (initial) (g) Weight 2 (final) (g) Histological cell scores Trichrome fibrosis scores
Yildirim Durak et al., 2013 [11]	Effect of vitamin C on the growth of experimentally induced endometriotic cysts	Female Wistar Albino Rats (n = 40) V1 group (n = 10) V2 group (n = 10) V3 group (n = 10) C group (n = 10)	6 weeks	Surgical induced endometriosis (autograft model) V1 group: 0.5 mg (2mg/kg) vit C/1 mL water for 6 weeks V2 group: 1.25 mg (5 mg/kg) vit C/1 mL water for 6 weeks V3 group: 2.5 mg (10 mg/kg) vit C/1 mL water for 6 weeks C group: distilled water 1 mL for 6 weeks Final surgical assessment	Vitamin C vs no vitamin C in all groups	Implant volume after operation (mm ³) Weight of cyst (mg) Histological cell scores Trichrome fibrosis scores NK cell contents
Eshrat Kalehoei et al., 2023 [13]	Therapeutic effects of L-arginine, L-carnitine, and mesenchymal stem cell-conditioned medium on endometriosis-induced oocyte poor quality in an experimental mouse model	Adult female NMRI mice (6-8 weeks old) n = not stated 1. Normal group 2. EMS-induced group		1. EMS induction 2. IVF LA: 250 mg/kg LC: 250 mg/kg LC BMSC-CM: 100 µL of CM/mouse	1. control 2. 250 mg/kg LA 3. 250 mg/kg LC 4. 100 µL BMSC-CM	1. <i>In vitro</i> maturation of immature oocytes: GV (%) GVBD (%) MII (%) DEG (%) 2. Blastocysts cell number N. Blast N. total cells N. TE N. ICM TE/ICM 3. The percentage of different steps of mice embryo development N. MII IVF (%) Cleavage (%) Morula (%) 4. Blood serum antioxidant capacity TAC (nmol/mL) NO (nmol/mL) TOS (nmol/mL) OSI All these outcomes are assessed in both EMS and normal groups and according to different antioxidant therapy (CO, LA, LC, BMSC-CM)
Hayedeh Hoorsan et al., 2022 [12]	The effectiveness of antioxidant therapy (vitamin C) in an experimentally induced mouse model of ovarian endometriosis	Mature, virgin female NMRI mice (n = 14) A group (n = 7) B group (n = 7)	Not stated	Surgical induced endometriosis (autograft model) 2 nd surgery (assessment of the endometriotic implants) A group: 50 mg/kg (0.5 mL) vit C every 2 days for 4 weeks B group: a 0.5 mL mix of water and starch Final surgical assessment	Vitamin C vs no vitamin C in all groups	Implant volume at the 2 nd operation (mm ³) Implant volume at the 3 rd operation (mm ³) Weight 1 (initial) (g) Weight 2 (final) (g) Histological cell scores Trichrome fibrosis scores Follicle number Atretic follicle number Corpus luteum number

NK: natural killer; vit C: vitamin C; EMS: endometriosis; IVF: *in vitro* fertilization; LA: L-arginine; LC: L-carnitine; BMSC-CM: bone morphogenetic stem cells; GV: Germinal vesicle; GVBD: Germinal vesicle break down; MII: metaphase II; DEG: degenerate oocytes; GVBD: Germinal vesicle break down; TOS: total oxidant status; NO: nitric oxide; TAC: total antioxidant capacity; OSI: oxidative stress index; CO: control.

Table 2. Description of selected studies included in the review (patients - human).

First author, year of publication	Title	Population	Follow up period	Intervention	Comparison	Outcomes
Vanessa SI et al., 2015 [15]	N-Acetyl-Cysteine and L-Carnitine Prevent Meiotic Oocyte Damage Induced by Follicular Fluid From Infertile Women With Mild Endometriosis	FF samples from infertile women (n = 22) 1. EMS-associated infertility (n = 11) 2. Other infertility (n = 11)	February 2009 - February 2011	1. Laparoscopic surgery in women with EMS. 2. FF-collection 3. Bovine oocyte collection 4. <i>In vitro</i> maturation 1. NAC 1.5 mmol/L 2. LC 0.6 mg/mL 3. NAC 1.5 mmol/L + LC 0.6 mg/mL	1. (No-FF) 2. (CFF) 3. (C + NAC 1.5 mmol/L) 4. (C + LC 0.6 mg/mL) 5. (C + 2Ao); 6. (EFF) 7. (E + NAC 1.5 mmol/L) 8. (E + LC 0.6 mg/mL) 9. (E + 2Ao).	MI, n (%) TI, n (%) PA, n (%) Total no. of MI, n (%) Analyzable MI, n (%) Normal MI, n (%)
Vanessa SI et al., 2021 [16]	Follicular Fluid from Infertile Women with Mild Endometriosis Impairs <i>In Vitro</i> Bovine Embryo Development: Potential Role of Oxidative Stress	FF samples from infertile women (n=22) 1. EMS-associated infertility (n = 11) 2. Other infertility (n = 11)	February 2009 - February 2011	1. Laparoscopic surgery in women with EMS. 2. FF-collection 3. Bovine oocyte collection 4. <i>In vitro</i> maturation 5. <i>In vitro</i> fertilization 6. <i>In vitro</i> embryo culture 1. NAC 1.5 mmol/L 2. LC 0.6 mg/mL 3. NAC 1.5 mmol/L + LC 0.6 mg/mL	1. (No-FF) 2. (CFF) 3. (C + NAC 1.5 mmol/L) 4. (C + LC 0.6 mg/mL) 5. (C + 2Ao) 6. (MEFF) 7. (MEFF + NAC 1.5 mmol/L) 8. (MEFF + LC 0.6 mg/mL) 9. (MEFF + 2Ao).	Presumptive zygotes (n) Cleavage rate % (n) Blastocysts formation rate % (n) Hatching rate % (n)
Xiang Lu et al., 2018 [14]	Effects of vitamin C on the outcome of <i>in vitro</i> fertilization-embryo transfer in endometriosis: A randomized controlled study	Patients with EMs (n = 280) Group 1 - Vit C treatment (n = 160) Group 2 - no vit C (n = 120) Patients with no EMs (n = 150)	June 2013 -December 2016.	1. IVF-ET procedure 2. Vit. C treatment Group 1 (n = 160) received 1000 mg/day from 2 months before IVF-ET treatment until 2 weeks after ET	EMS patients vs no EMS patients EMS patients treated with vit C/ not treated with vit C	1. Laboratory and pregnancy outcomes in EMS patients/ no EMS patients Total Gn dosage No. of retrieved oocytes Fertilization rate (%) High-grade embryo rate (%) Implantation rate (%) Clinical pregnancy rate (%) No. of frozen embryos 2. Changes in serum levels of vit C and oxidative stress markers in EMS patients with vit C/ no vit C Serum levels of vit C (µmol/L) Serum levels of SOD (U/L) Serum levels of TAC (mmol/L) Serum levels of MDA (µM) Serum levels of ROS (CPS)
Jennifer Mier-Cabrera et al., 2008 [17]	Effect of vitamins C and E supplementation on peripheral oxidative stress markers and pregnancy rate in women with endometriosis	Patients with EMs (n = 34) Group 1 - Vit C and Vit E treatment (n = 16) Group 2 - placebo (n = 18)	15 months	Group 1 - 343 mg of vitamin C and 84 mg of vitamin E	Vit C and vit E patients group vs placebo group	1. Oxidative stress marker levels in women with endometriosis (Baseline, at 2 months, at 4 months, at 6 months in PF, plasma) LOOH (µmol/L) MDA (µmol/L) 2. Pregnancy rate

First author, year of publication	Title	Population	Follow up period	Intervention	Comparison	Outcomes
Nalini Santanam <i>et al.</i> , 2016 [18]	Myeloperoxidase as a Potential Target in Women With Endometriosis Undergoing IV	Patients (n = 117) Complete data (n = 68) No EMs group (n = 41) Mild EMs group (n = 20) Moderate/severe EMs group (n = 7)	Not stated	1. IVF 2. Collection of FF 3. Collection of Blood Plasma Patients received 800 IU of vit E and 1000 IU of vit C for a minimum of 8 weeks No EMs group vit C+E (n = 5) Mild EMs group vit C+E (n = 5) Moderate/severe EMs group vit C+E (n = 4)	Vit C and vit E patients group vs placebo group	MPO level (ng/ml)
Sahar Rostami <i>et al.</i> , 2023 [20]	Astaxanthin ameliorates inflammation, oxidative stress, and reproductive outcomes in endometriosis patients undergoing assisted reproduction: A randomized, triple-blind placebo-controlled clinical trial	Infertile patients (n = 73) with EMs Complete data (n = 50) AST group (n = 25) Placebo group (n = 25)	December 2021 - September 2022.	1. IVF and FF collection 2. Blood and FF collection AST group: 6 mg daily of oral AST for 12 weeks Placebo group: 6 mg daily of placebo capsules for 12 weeks	AST treatment vs placebo	1. OS markers and cytokine levels MDA SOD CAT TAC L-1b IL-6 TNF-a 2. ART outcomes Number of oocytes GV MI MII Oocyte maturity rate (MI (%)) Fertilized Fertilization rate (%) Number of frozen embryos High-quality embryos Frozen embryos Number of transferred embryos
Leila Amimi <i>et al.</i> , 2021 [19]	The Effect of Combined Vitamin C and Vitamin E Supplementation on Oxidative Stress Markers in Women with Endometriosis: A Randomized, Triple-Blind Placebo-Controlled Clinical Trial	Patients with endometriosis (n = 60) A group (n = 30) B group (n = 30)	June 2017-November 2017	A group: vitamin C 1000 mg/day (2 tablets/500 mg) + vitamin E 800 IU/day (2 tablets/400 IU) for 8 weeks. B group (placebo pills (mannitol and magnesium stearate polyvinylpyrrolidone)) for 8 weeks	Vit C and vit E patients group vs placebo group	1. OS markers levels MDA ROS TAC dyspareunia 2. VAS score of dysmenorrhea,

CFF: control follicular fluid; OS: oxidative stress; ROS: reactive oxygen species; LC: L-carnitine; NAC: N-Acetyl-Cysteine; BMSC-CMI: bone morphogenetic stem cells; TOS: total oxidant status; NO: nitric oxide; TAC: total antioxidant capacity; OSI: oxidative stress index; IVF: *In vitro* fertilization; SOD: superoxide dismutase; MDA: malondialdehyde; FF: follicular fluid; EFF: endometriosis follicular fluid; LOOH: lipid hydroperoxide; MPO: myeloperoxidase; AST: astaxanthin; ART: associated reproductive technology; CAT: catalase; TAR: Total antioxidant response; NK: Natural killer; EMS: endometriosis; TNF-a: Tumor necrosis factor; IL-1b: interleukin 1b; IL-6: interleukin 6; GV: germinal vesicle; MI: metaphase I; PA: spontaneous parthenogenetic activation; TI: telophase.

dies, the authors found a significant difference not only in the volume of endometriotic implants but also in the trichrome fibrosis scores ($p = 0.03$). Follicle, atretic follicle and corpus luteum counts were also measured. The number of follicles was significantly increased ($p = 0.0005$) and the number of atretic follicles was significantly decreased ($p = 0.006$) after vitamin C therapy.

L-arginine, L-carnitine effectiveness in animal population

The other study based on the induced endometriosis mouse model was conducted by Eshrat Kalehoei *et al.* [13]. The authors compared the effects of L-arginine (LA), L-carnitine (LC), bone morphogenetic stem cells (BMSC-CM) on endometriosis-induced oocyte quality and levels of oxidative stress markers. In the endometriosis group, mice treated with LC, LA or BMSC-CM had significantly lower levels of oxidative stress markers compared to control. In normal and endometriosis treatment groups *in vitro* fertilisation (IVF), cleavage and blastocyst formation rates were significantly improved compared to the control ($p < 0.05$).

Antioxidants effectiveness in human population

Other 7 studies were based on the follicular fluid and plasma samples collected from women with endometriosis related infertility.

The vitamin C effectiveness in human population

In human population, the efficacy of vitamin C supplementation was analysed by Lu *et al.* [14]. There was no significant difference in fertilization, implantation and pregnancy rates between all participants. The number of oocytes and frozen embryos in endometriosis groups was significantly lower than in control ($p < 0.05$). Treatment with vitamin C for 2 months improved its serum and follicular fluid concentration in patients with endometriosis, however, it did not affect oxidative stress markers rate. The results could be as follows because of several limitations: small sample size and only one time point measurement (2 months of vitamin C supplementation): comparison of different time points was not performed.

N-acetyl-cysteine, L-carnitine effectiveness in human population

Giorgi *et al.* conducted two studies [15, 16] in which the percentage of meiotically normal oocytes in metaphase II, presumptive zygotes, cleavage rate, blastocyst formation rate and hatching rate were

measured. The number of meiotically abnormal and normal metaphase II oocytes was similar in all 9 groups. The authors suggest that follicular fluid (FF) from infertile women with endometriosis increases the percentage of meiotically abnormal oocytes. There was no significant difference between groups in cleavage ($p = 0.54$) and blastocyst formation ($p = 0.4349$) rates. However, the hatching rate was higher in the control follicular fluid group than in the endometriosis follicular fluid (EFF) group. The addition of antioxidants in CFF groups did not affect the hatching rate. The addition of N-acetylcysteine reduces the destructive effects of FF on the oocyte meiotic spindle and increases hatching rate. The addition of L-carnitine completely prevents this destructive effect on the meiotic spindle, but is less effective in terms of hatching rate.

The vitamin C and vitamin E combination effectiveness in human population

The effect of combined vitamin C + vitamin E treatment was evaluated by Mier-Cabrera *et al.* [17], Santanam *et al.* [18] and Amini *et al.* [19].

Mier-Cabrera *et al.* [17] analysed the lipid hydroperoxide (LOOH) and malondialdehyde (MDA) rate in plasma and peritoneal fluid. There was a statistically significant difference in plasma LOOH and MDA concentrations between control and treatment groups. Amini *et al.* [19] held randomized, triple-blind placebo-controlled clinical trial, where statistically reduced MDA ($p = 0.002$) and reactive oxygen species (ROS) ($p < 0.001$) levels in treatment group compared to placebo were evaluated.

Anti-myeloperoxidase therapy human population

Santanam *et al.* [18] suggested that the level of myeloperoxidase (MPO) – one of the oxidative stress markers – depends on the severity of endometriosis. Mean MPO levels in follicular fluid collected from women with severe endometriosis were significantly higher than in control and mild endometriosis groups. Combination antioxidant treatment did not significantly reduce MPO levels in both groups.

Astaxanthin therapy in human population

Rostami *et al.* conducted a study on the efficacy of astaxanthin (AST) on oxidative stress markers, cytokine levels and associated reproductive technology (ART) outcomes in infertile women with endo-

metriosis [20]. All parameters were reduced after antioxidant therapy except serum catalase (CAT), IL-1b, IL-6 levels. Embryo quality, number of metaphase II oocytes improved significantly after therapy. However, the number of embryos transferred, fertilisation rate and pregnancy rate were similar in both groups.

Visualisation tools were provided by the ROBVIS application [21]. According to the ROBINS-I tool, the overall risk of bias for non-randomised trials was 66.7 % low and 33.3% serious (Figure 2). Based on the RoB 2 tool (Figure 3), randomized trials had a 75% chance of low risk of bias and an 25% chance of some concern regarding the overall risk of bias. The SYRCLE’s RoB tool was used to assess the quality of included animal studies (Figure 4). The risk of allocation concealment and random housing could not be confirmed because none of the studies offered complete information.

DISCUSSION

There is increasing evidence to suggest that specific diet patterns and nutrients may modulate the pathophysiological processes underlying endometriosis.

In this systematic review, we evaluated the efficiency of antioxidants supplementation in endometriosis treatment. We found out that antioxidants reduce the severity of endometriosis symptoms by affecting the pathogenesis of the disease.

Oxidative stress occurs when the balance between reactive oxygen species production and antioxidant capacity is disturbed, either by insufficient antioxidant protection or by increased ROS production. The relationship between ROS production and the progression of endometriosis has been studied previously [22]. Due to dysregulation of iron metabolism, these abnormal endometriotic lesions are




Figure 2. ROBINS-1 tool for non-randomized trials. Domains: D1: Bias due to confounding. - D2: Bias due to selection of participants. - D3: Bias in classification of interventions. - D4: Bias due to deviations from intended interventions. D5: Bias due to missing data. - D6: Bias in measurement of outcomes. - D7: Bias in selection of the reported result.




Figure 3. RoB2 tool for randomized trials. Domains: D1: Bias arising from the randomization process. - D2: Bias due to deviations from intended intervention. - D3: Bias due to missing outcome data. - D4: Bias in measurement of the outcome. - D5: Bias in selection of the reported result

	Sequence generation	Baseline characteristics	Allocation concealment	Random housing	Blinding (study team)	Random outcome assessment	Blinding (outcome assessors)	Outcome data	Selective outcome reporting	Other bias	
	Selection bias			Performance bias		Detection bias		Attrition bias	Reporting bias	Other bias	
Ozlem Ulas Erten et al (2016)	?	+	?	?	+	?	+	+	+	+	+
Yildirim Durak et al (2013)	?	+	?	?	+	?	+	+	+	+	+
Hayedeh Hoorsan et al (2022)	?	+	?	?	+	?	+	+	+	+	+
Eshrat Kalehoei et al (2023)	?	?	?	?	?	?	?	-	+	-	




 - Low risk
 - High risk
 - Unclear risk

Figure 4. SYRCLÉ’s risk of bias tool for each experiment al animal study.

thought to be resistant to ferroptosis. Ferroptosis is a form of regulated, iron-catalysed cell death caused by excessive lipid peroxidation in cell membranes. This process was first described by Dixon in 2012 [23].

Li *et al.* [24] found out that there was an excess expression of ferroptosis-associated genes in the ectopic and eutopic endometrium in patients with endometriosis, showing a general trend towards inhibition of the ferroptosis pathway. Increased transferrin receptors (TFR1) and Ras gene mutations in abnormal endometriotic cells directly affect ferroptosis resistance. A local imbalance in iron homeostasis leads to oxidative stress in the intraperitoneal cavity, inflammation and ferroptosis in intact peripheral tissues. Iron-dependent ROS synthesis is based on the Fenton reaction: $Fe^{2+} + H_2O_2 \rightarrow Fe^{3+} + OH^- + OH^-$. As a result, a hydroxyl radical (-OH) is formed, leading to lipid peroxidation and accumulation of lipid LOOH, which damages the membrane. This is why ectopic endometriotic tissue has higher levels of lipid peroxidation products than normal endometrial tissue. This is also confirmed by other studies [17, 25].

In addition, we observe changes in enzyme levels: superoxide dismutase (SOD) and indicators such as TAC, total oxidant status (TOS) and oxidative stress index (OSI) in serum and FF. The decrease in TAC and SOD between patients with and without endometriosis is confirmed. It is noted that there was a significant difference in this indicator in FF as opposed to serum between both groups [14]. Total antioxidant response (TAR) is also lower in patients with endometriosis, leading to an excess of OSI [26]. Treatment with AST improved TAC and SOD levels [19]. LC and LA administration also improved the TAC, reduced TOS, nitric oxide (NO) and OSI ($p < 0.05$) [16]. However, vitamin C treatment

showed no difference in oxidative stress markers and enzyme levels [14]. But there is evidence that vitamin C prevents the progression of endometriotic lesion development by reducing their weight, size and volume [10-12]. Vitamin C and vitamin E combination significantly suppressed levels of MPO (a neutrophil marker that is increased due to oxidative stress and depends on the severity of endometriosis) in FF [18]. These findings support previous data [26].

It should be mentioned that immune cells play a crucial role in ectopic endometriotic lesions detection and elimination. It is known that oxidative stress impairs the efficiency of the immune system, leading to reduced recognition of abnormal endometrial tissue, allowing its invasion, accumulation and growth in the pelvic and abdominal cavity. Antioxidants are known to stimulate the whole process of phagocytosis [27]. Durak *et al.* [11] found that NK cell (Natural killer cells) levels were significantly lower in control groups than in those on antioxidant therapy ($p < 0.01$). Similar results have been reported in other studies [28, 29]. It should be noted that the decrease in cellular immunity correlates with the severity of the disease. Whether this decrease in NK cells is a cause or a consequence of the severity of endometriosis remains unclear. The reduction in cellular immunity is related to the “endometriotic disease theory”, also known as Sampson’s theory, according to which the most important factor in the development of endometriosis is not the initial implantation in the peritoneal cavity, but cell mutations that cannot be eliminated due to the reduced number of immune cells. It is these ectopic endometriotic cells, ignored by regulatory factors in the peritoneal fluid, that trigger the disease [30].

All of the above factors affect fertility in women with endometriosis. The granulosa cells and the surrounding cumulus cells in the follicle are involved in the maturation of the oocyte. This process depends on the intrafollicular environment. If it is damaged, the developmental competence of the oocytes, the quality of the embryos and the clinical pregnancy rate are reduced. In the case of antioxidant therapy, IVF, cleavage and blastocyst formation rates are increased compared to no treatment. It is also important to highlight that this finding is potentially helpful for translation into clinical practice [13-16, 19]. High-level antioxidant diet can significantly influence inflammatory processes, which are directly related to the pathophysiology of endometriosis.

Strengths and limitations of the study

The limitations we encountered were mainly related to the available data sources. The patients were not similar between studies: rats and mice with induced endometriosis, human. The heterogeneity of the antioxidants should also be mentioned. A total of 6 antioxidants were included in this review, but it is difficult to compare them because of differences in regimen and dosage in each study. Studies in animal models have a lower quality of evidence than those in humans. It is important to emphasize that more research is needed in human to assess the clinical relevance and to establish the efficacy of antioxidants, as clinical trials evaluating their effects on endometriosis are still relatively limited. The main strength of this study is that we observed antioxidant supplementation as a therapy that affect oxidative stress – the main aspect of endometriosis pathogenesis. All previously published reviews were aimed to analyse the types of oxidative stress markers and their levels in patients with endometriosis, but did not observe and summarize any medications for their reduction.

Implications for future studies may include investigating the development of targeted antioxidant treatment, the possibility of delivering antioxidants directly to endometriotic lesions. This could potentially increase the efficacy of antioxidant therapy and minimize potential side effects.

CONCLUSIONS

Antioxidants may be considered as a possible component of endometriosis therapy to potentially

enhance fertility outcomes and slow disease progression, though current evidence is preliminary and requires further validation. This type of treatment reduces oxidative stress markers concentration, suspend endometriotic lesions progression and improve oocyte developmental competence. However, there is still controversy about the antioxidant treatment as a monotherapy of endometriosis. That is why more clinical trials to make stronger recommendations is needed.

COMPLIANCE WITH ETHICAL STANDARDS

Authors' contribution

L.P., E.N., S.I., V.T., A.L., V.S.: Conceptualization. E.N., I.S., A.L.: Data curation, writing – original draft. L.P., V.T., V.S.: Writing – review & editing.

Funding

None.

Study registration

PROSPERO CRD42023454705.

Disclosure of interests

The authors declare that they have no conflict of interests.

Ethical approval

N/A.

Informed consent

N/A.

Data sharing

N/A.

REFERENCES

1. Czyzyk A, Podfigurna A, Szeliga A, Meczekalski B. Update on endometriosis pathogenesis. *Minerva Ginecol.* 2017;69(5):447-461. doi: 10.23736/S0026-4784.17.04048-5.
2. Ng SW, Norwitz SG, Taylor HS, Norwitz ER. Endometriosis: The role of iron overload and ferroptosis. *Reprod Sci.* 2020;27(7):1383-1390. doi: 10.1007/s43032-020-00164-z.
3. Kobayashi H, Yoshimoto C, Matsubara S, Shigetomi H, Imanaka S. Current understanding of and future directions for endometriosis-related

- infertility research with a focus on ferroptosis. *Diagnostics* (Basel). 2023;13(11):1926. doi: 10.3390/diagnostics13111926.
4. Kao SH, Huang HC, Hsieh RH, Chen SC, Tsai MC, Tzeng CR. Oxidative damage and mitochondrial DNA mutations with endometriosis. *Ann NY Acad Sci*. 2005;1042:186-194. doi: 10.1196/annals.1338.021.
 5. Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ*. 2021;372:n160. doi:10.1136/bmj.n160.
 6. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. *Cochrane Handbook for Systematic Reviews of Interventions* version 6.1 (updated September 2020). Cochrane, 2020. Available from: www.training.cochrane.org/handbook.
 7. Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: a revised tool for assessing risk of bias in randomized trials. *BMJ*. 2019;366:l4898. doi: 10.1136/bmj.l4898.
 8. Sterne JAC, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M, et al. ROBINS-I: a tool for assessing risk of bias in nonrandomized studies of interventions. *BMJ*. 2016;355:i4919. doi: 10.1136/bmj.i4919.
 9. Hooijmans CR, Rovers MM, de Vries RB, Lee-naars M, Ritskes-Hoitinga M, Langendam MW. SYRCLE's risk of bias tool for animal studies. *BMC Med Res Methodol*. 2014;14:43. doi: 10.1186/1471-2288-14-43.
 10. Erten OU, Ensari TA, Dilbaz B, Cakiroglu H, Altinbas SK, Çaydere M, et al. Vitamin C is effective for the prevention and regression of endometriotic implants in an experimentally induced rat model of endometriosis. *Taiwan J Obstet Gynecol*. 2016;55(2):251-257. doi: 10.1016/j.tjgo.2015.07.004.
 11. Durak Y, Kokcu A, Kefeli M, Bildircin D, Çelik H, Alper T. Effect of vitamin C on the growth of experimentally induced endometriotic cysts. *J Obstet Gynaecol Res*. 2013;39(7):1253-1258. doi: 10.1111/jog.12050.
 12. Hoorsan H, Simbar M, Tehrani FR, Fathi F, Mosaffa N, Riazi H, et al. The effectiveness of antioxidant therapy (vitamin C) in an experimentally induced mouse model of ovarian endometriosis. *Womens Health (Lond)*. 2022;18:17455057221096218. doi: 10.1177/17455057221096218.
 13. Kalehoei E, Moradi M, Azadbakht M, Zhaleh H, Abadi SAL, Mahdiuni H, et al. Therapeutic effects of L-arginine, L-carnitine, and mesenchymal stem cell-conditioned medium on endometriosis-induced oocyte poor quality in an experimental mouse model. *J Obstet Gynaecol Res*. 2023;49(4):1180-1188. doi: 10.1111/jog.15569.
 14. Lu X, Wu Z, Wang M, Cheng W. Effects of vitamin C on the outcome of in vitro fertilization-embryo transfer in endometriosis: A randomized controlled study. *J Int Med Res*. 2018;46(11):4624-4633. doi: 10.1177/0300060518786918.
 15. Giorgi VS, Da Broi MG, Paz CC, Ferriani RA, Navarro PA. N-acetyl-cysteine and l-carnitine prevent meiotic oocyte damage induced by follicular fluid from infertile women with mild endometriosis. *Reprod Sci*. 2016;23(3):342-351. doi: 10.1177/1933719115602772.
 16. Giorgi VSI, Ferriani RA, Navarro PA. Follicular fluid from infertile women with mild endometriosis impairs in vitro bovine embryo development: Potential role of oxidative stress. *Rev Bras Ginecol Obstet*. 2021;43(2):119-125. doi: 10.1055/s-0040-1718443.
 17. Mier-Cabrera J, Genera-García M, De la Jara-Díaz J, Perichart-Perera O, Vadillo-Ortega F, Hernández-Guerrero C. Effect of vitamins C and E supplementation on peripheral oxidative stress markers and pregnancy rate in women with endometriosis. *Int J Gynaecol Obstet*. 2008;100(3):252-256. doi: 10.1016/j.ijgo.2007.08.018.
 18. Santanam N, Zoneraich N, Parthasarathy S. Myeloperoxidase as a potential target in women with endometriosis undergoing IVF. *Reprod Sci*. 2017;24(4):619-626. doi: 10.1177/1933719116667225.
 19. Amini L, Chekini R, Nateghi MR, Haghani H, Jamialahmadi T, Sathyapalan T, et al. The effect of combined vitamin C and vitamin E supplementation on oxidative stress markers in women with endometriosis: A randomized, triple-blind placebo-controlled clinical trial. *Pain Res Manag*. 2021;2021:5529741. doi: 10.1155/2021/5529741.
 20. Rostami S, Alyasin A, Saedi M, Nekoonaam S, Khodarahmian M, Moeini A, et al. Astaxanthin ameliorates inflammation, oxidative stress, and reproductive outcomes in endometriosis patients undergoing assisted reproduction: A randomized, triple-blind placebo-controlled clinical trial. *Front Endocrinol*. 2023;14:1144323. doi: 10.3389/fendo.2023.1144323.

21. McGuinness LA, Higgins JPT. Risk-of-bias VI-Sualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments. *Res Syn Meth*. 2020;12(1):55-617. doi: 10.1002/jrsm.1411.
22. Scutiero G, Iannone P, Bernardi G, Bonaccorsi G, Spadaro SA, Volta CA, et al. Oxidative stress and endometriosis: A systematic review of the literature. *Oxid Med Cell Longev*. 2017;2017:7265238. doi: 10.1155/2017/7265238.
23. Dixon SJ, Lemberg KM, Lamprecht MR, Skouta R, Zaitsev EM, Gleason CE, et al. Ferroptosis: An iron-dependent form of nonapoptotic cell death. *Cell*. 2012;149(5):1060-1072. doi: 10.1016/j.cell.2012.03.042.
24. Li B, Duan H, Wang S, Li Y. Ferroptosis resistance mechanisms in endometriosis for diagnostic model establishment. *Reprod Biomed Online*. 2021;43(1):127-138. doi: 10.1016/j.rbmo.2021.04.002.
25. de Lima CB, Cordeiro FB, Camargo M, Zylbersztejn DS, Cedenho AP, Bertolla RP, et al. Follicular fluid lipid peroxidation levels in women with endometriosis during controlled ovarian hyperstimulation. *Hum Fertil (Camb)*. 2017;20(1):48-54. doi: 10.1080/14647273.2016.1246753.
26. ODF, Waelkens E, Peterse DP, Lebovic D, Meuleman C, Tomassetti C, et al. Evaluation of total, active, and specific myeloperoxidase levels in women with and without endometriosis. *Gynecol Obstet Investig*. 2018;83(2):133-139. doi: 10.1159/000475664.
27. Assaf L, Eid AA, Nassif J. Role of AMPK/mTOR, mitochondria, and ROS in the pathogenesis of endometriosis. *Life Sci*. 2022;306:120805. doi: 10.1016/j.lfs.2022.120805.
28. Fan D, Wang X, Shi Z, Jiang Y, Zheng B, Xu L, et al. Understanding endometriosis from an immunomicroenvironmental perspective. *Chin Med J (Engl)*. 2023;136(16):1897-1909. doi: 10.1097/CM9.0000000000002649.
29. Wang L, Li L, Li Y, Huang C, Lian R, Wu T, et al. History of endometriosis is associated with decreased peripheral NK cytotoxicity and increased infiltration of uterine CD68+ macrophages. *Front Immunol*. 2021;12:711231. doi: 10.3389/fimmu.2021.711231.
30. Chen S, Liu Y, Zhong Z, Wei C, Liu Y, Zhu X. Peritoneal immune microenvironment of endometriosis: Role and therapeutic perspectives. *Front Immunol*. 2023;14:1134663. doi: 10.3389/fimmu.2023.1134663.