



Italian Journal of
Gynæcology & Obstetrics
September 2026 - Vol. 38 - N. 3 - Quarterly - ISSN 2385 – 0868

Comparative efficacy of letrozole *versus* letrozole-metformin in the ovulation and clinical pregnancy rate in polycystic ovary syndrome: a prospective randomized controlled study

Sakshi Chopra *, Ruchi Hooda, M. Gouri Devi

Department of Reproductive Medicine, Ridge IVF Centre, New Delhi, India.

ARTICLE INFO

History

Received: 02 August 2025

Received in revised form: 22 November 2025

Accepted: 22 January 2026

Available online: 10 September 2026

DOI: 10.36129/jog.2026.261

Key words

PCOS; letrozole; metformin; ovulation.

***Corresponding author:** Sakshi Chopra, MBBS
DGO DNB FRM. Department of Reproductive
Medicine, Ridge IVF Centre, 109, United India
Apts, Mayur Vihar, Phase 1, extension, Delhi
110091, India.

Email: 385saki@gmail.com.

ORCID: 0000-0002-4905-2237.

ABSTRACT

Objective. The study provides an insight on the effect of letrozole and letrozole-metformin combination on the formation of mature follicle, ovulation, endometrial thickness and pregnancy rates in anovulatory polycystic ovary syndrome (PCOS) infertile women.

Materials and Methods. A prospective randomised controlled study conducted at Ridge IVF Centre, Delhi from May 2024-April 2025. 120 women of ages 21-40 years affected by PCOS and trying naturally for conception were included. Patients were randomly selected into 2 groups: Group A (letrozole and metformin) and Group B (letrozole only).

Metformin 500 mg twice daily was added from day 1 of cycle. Letrozole 2.5 mg BD was started from day 2/3 of menstrual cycle. Follicular monitoring was done from day 10 by transvaginal sonography and once a mature follicle of size > 18 mm is seen on TVS, injection. hCG 10,000 IU was given. Once ovulation was documented, couple was advised for alternate day intercourse and luteal phase support was given for 14 days. Pregnancy confirmation was done by S.βhCG after 14 days.

Results. The study showed that the formation of mature follicle was more in letrozole-metformin group and ovulation rate was higher in letrozole only group. The endometrial thickness on the day of the trigger was comparable in both groups. The clinical pregnancy was slightly higher in combination group, but not statistically significant.

Conclusions. Adding metformin to letrozole does insignificant improvement in ovulation patterns in PCOS anovulatory women.

INTRODUCTION

Polycystic ovary syndrome (PCOS) is the most common endocrinological disorder affecting 4-12% of women [1-2]. It has been the most controversial

medical condition affecting women of reproductive age group. It is one of the causes for anovulatory female infertility. PCOS is a heterogeneous disorder with a combination of genetic, reproductive and metabolic characteristics. These women are at risk

of Type 2 diabetes mellitus (DM) and cardiovascular disease.

According to 2003 Rotterdam Consensus revised diagnostic criteria, at least two of the following is required for the diagnosis of PCOS:

1. oligo or anovulation, or both, that is, menstrual disturbance;
2. clinical or biochemical signs, or both, of hyperandrogenism;
3. PCO on ultrasound and exclusion of other aetiologies of menstrual disturbance and hyperandrogenism (such as congenital adrenal hyperplasia, androgen-secreting tumours, Cushing's syndrome) [3].

A recent update to guidelines in view of advancement in ultrasound technology and resolution, state that the diagnostic criteria for ultrasound PCO morphology is either 20 or more follicles per ovary or increased ovarian volume, over 10 mL, when using a transvaginal ultrasound scan [4].

Insulin resistance is a very common feature of PCOS, owing to either genetic propensity or obesity [5]. It is not a diagnostic feature of this disorder. In PCOS, hyperinsulinemia can induce androgen production and decrease the levels of sex hormone binding globulin. A high level of androgen levels affects the oocytes growth and causes atresia of follicles. PCOS is responsible for about 80% of anovulatory infertility [6]. Women with insulin resistance have increased prevalence of hirsutism than women with non-insulin-resistant PCOS [7]. These women are more likely to develop resistance to ovulation induction treatment.

Lifestyle modification including weight loss and exercise reduces central fat, improves insulin sensitivity, reduces testosterone levels and restoring ovulation in overweight, infertile women with PCOS [7].

Metformin is the member of the biguanide family that has been used for the treatment of T2DM. Metformin works by improving the insulin sensitivity of peripheral tissues [8], which results in a reduction of circulating insulin levels. The main side effects associated with metformin treatment are the gastrointestinal symptoms of nausea, diarrhoea, flatulence, bloating, anorexia, metallic taste and abdominal pain, causing reluctance in its use over a prolonged period [9].

Within the ovary, metformin has a direct impact on cells to reduce excessive steroidogenesis and folliculogenesis [9]. It reduces theca cell proliferation, decreases androgen production, reduces pituitary

luteinizing hormone and increase sex hormone binding globulin by the liver and promotes ovulation. Letrozole is a nonsteroidal reversible and competitive aromatase inhibitor. Letrozole has a half-life of about 45 hours [10]. It has a central effect on the pituitary-hypothalamic axis by releasing it from estrogen negative feedback and a local ovarian effect by blocking androgen conversion to estrogen, with the concomitant accumulation of androgens inside the ovary, augmenting the follicular FSH receptor expression, and promoting folliculogenesis [11]. This study provides an insight on the effect of letrozole and letrozole-metformin combination on anovulatory PCOS infertile women in ovulation and pregnancy.

MATERIALS AND METHODS

Study site

The study was a prospective randomised controlled study conducted in Department of Assisted Reproduction at Ridge IVF centre, New Delhi from May 2024-April 2025. The approval was taken from the ethics committee on board for the study.

Inclusion criteria

Women of all ages (21-40 years) who were affected by PCOS according to Rotterdam criteria with normal hysterosalpingogram and were trying naturally for conception were included.

Exclusion criteria

Patients with hormonal disorder like hyperprolactinemia, hypo or hyperthyroidism or diabetes, endometriosis, uterine fibroid, male factor infertility were excluded from study.

All patients were admitted through OPD after complete evaluation with history, examination, investigation and verbal informed consent for inclusion in the study. Blood for Insulin Postprandial is sent 1 ½ hours after meals on first visit along with other investigations.

Patients were selected randomly into 2 groups: Group A (letrozole and metformin) and Group B (letrozole only). Letrozole and metformin was administered as oral preparations.

In Group A, tablet metformin 500 mg twice daily with meals starting from day 1 of cycle was added and continued for three months and continued even after conception till first 3 months of pregnancy. Patients were followed for 3 months to evaluate

for pregnancy in both the groups. Oral ovulogen letrozole 2.5 mg BD was started from day 2/3 of menstrual cycle after spontaneous or withdrawal bleeding in both groups. Follicular monitoring was done from day 10 and follicle growth was monitored by transvaginal sonography (TVS). Once a mature follicle of size > 18 mm was seen, injection hCG 10,000 IU was given as a trigger. Once follicular rupture was documented, couple was advised for alternate day intercourse and luteal phase support was given with dydrogesterone 10 mg BD for 14 days. Pregnancy confirmation was done by estimating the S.βhCG concentration in the blood after 14 days and doubling of the S.βhCG after 48 hours. The clinical pregnancy was established by the presence of gestational sac by ultrasonography performed at 6-7 weeks of pregnancy.

Treatment was repeated for 3 months if the patient fails to ovulate or achieve pregnancy.

The primary outcomes were assessed as documentation of mature follicle (> 18mm) and documentation of ovulation (ovulation rate).

The secondary outcomes were assessed as measurement of ovarian volume, hormonal profile on day 2 (S.FSH, S.LH, S.E2), endometrial thickness on day of hCG trigger and clinical pregnancy rate.

Statistical analysis

The collected data was entered in Microsoft Excel and then analysed and statistically evaluated using SPSS-25 version. Normality of each variable was assessed by using the Kolmogorov-Smirnov test. Quantitative data was expressed by mean, standard deviation and difference between means of two group were tested by Unpaired t test or Mann-Whitney U test. Qualitative data were expressed in percentage and difference between percentage of two group were tested by chi square test or Fisher exact test. P-value less than 0.05 was considered statistically significant.

RESULTS

The baseline characteristics of participants in both groups were largely comparable (**Table 1**). The mean age was similar in Group A (30.13 ± 3.21 years) and Group B (30.22 ± 3.25 years). The mean BMI was slightly higher in Group A (27.68 ± 4.78 kg/m²) compared to Group B (26.22 ± 4.12 kg/m²), though this difference was not statistically significant ($p = 0.07$). In terms of type of infertility, pri-

Table 1. Data comparing the baseline characteristics between Group A and Group B.

	Group A (letrozole-metformin) (n = 60)	Group B (letrozole) (n = 60)	P-value
Mean age in years	30.13 ± 3.21	30.22 ± 3.25	0.88
Mean BMI (kg/m ²)	27.68 ± 4.78	26.22 ± 4.12	0.07
Type of infertility			
Primary	46(76.7%)	48(80.0%)	0.65
Secondary	14(23.3%)	12(20.0%)	
Cycle			
Irregular	34 (56.7%)	11 (18.3%)	<0.001
Regular	26 (43.3%)	49 (81.7%)	<0.001

BMI: body mass index; n: number of participants. Data are presented as mean $\pm$ standard deviation (SD) or n %.

Table 2. Data comparing the other parameters between Group A and Group

	Group A (letrozole-metformin) (n = 60)	Group B (letrozole) (n = 60)	P-value
RT ovary VOL (ml)	9.83 ± 1.48	6.76 ± 1.94	<0.001
LT ovary VOL (ml)	9.24 ± 1.07	6.86 ± 2.01	<0.001
S.AMH (ng/ml)	8.12 ± 3.42	6.13 ± 2.44	0.001
S.FSH	6.50 ± 1.29	6.02 ± 1.61	0.05
S.LH	7.40 ± 3.19	6.31 ± 4.17	0.01
S.E2	43.27 ± 17.42	36.65 ± 13.07	0.02
Insulin PP (μU/ml)	67.98 ± 27.23	26.92 ± 15.62	<0.001
Endometrial thickness (mm)	8.83 ± 1.65	8.80 ± 1.21	0.92

RT: right; LT: left; VOL: volume; AMH: anti mullerian hormone; LH: luteinizing hormone; E2: estradiol; PP: postprandial; n: number of participants. Data are presented as mean $\pm$ standard deviation (SD).

mary infertility was predominant in both groups: 76.7% in Group A and 80.0% in Group B.

However, a significant difference was observed in menstrual cycle regularity. 56.7% of Group A had irregular cycles compared to only 18.3% in Group B, and regular cycles were more common in Group B with statistical difference < 0.001 (**Table 1**).

In **Table 2**, comparison of hormonal and ultrasonographic parameters were done. Right ovarian volume was significantly higher in Group A (9.83 ± 1.48 ml) compared to Group B (6.76 ± 1.94 ml), and similarly, the left ovarian volume was also greater in Group A (9.24 ± 1.07 ml vs 6.86 ± 2.01 ml) with statistically significant ($p < 0.001$) (**Figure 1**).

Serum AMH levels were high in Group A (8.12 ± 3.42 ng/ml) as compared to Group B (6.13 ± 2.44 ng/ml). Baseline Serum FSH was marginally higher in Group A (6.50 ± 1.29) than in Group B (6.02 ± 1.61), with borderline significance ($p = 0.05$). Serum LH levels were significantly hi-




Figure 1. Ovarian volume of right and left ovary.




Figure 2. Serum LH on day 2 of cycle




Figure 3. Insulin post prandial.

gher in Group A (7.40 ± 3.19) compared to Group B (6.31 ± 4.17) ($p = 0.01$) (Figure 2) and serum estradiol (E2) levels were also higher in Group A (43.27 ± 17.42 vs 36.65 ± 13.07).

A striking difference was observed in postprandial insulin levels, which were markedly elevated in Group A (67.98 ± 27.23 μ U/ml) compared to Group B (26.92 ± 15.62 μ U/ml), showing a highly significant difference ($p < 0.001$) (Figure 3).

Endometrial thickness on the trigger day was comparable between both groups (8.83 ± 1.65 mm in Group A vs 8.80 ± 1.21 mm in Group B (Figure 4).

Table 3. Data comparing clinical hirsutism between Group A and Group B.

Clinical hirsutism	Group A (letrozole-metformin) (n = 60)	Group B (letrozole) (n = 60)	P-value
Absent	35 (58.3%)	50 (83.3%)	<0.01
Present	25 (41.7%)	10 (16.7%)	

Table 4. Data comparing different outcome between Group A and Group B.

	Group A (letrozole-metformin) (n = 60)	Group B (letrozole) (n = 60)	P-value
Mature follicle			0.01
No	7(11.7%)	0	
Yes	53(88.3%)	60	
Ovulation			0.002
No	18(30.0%)	4(6.7%)	
Yes	42(70.0%)	56(93.3%)	
S.βhCG			0.07
Negative	36(60.0%)	45(75.0%)	
Positive	24(40.0%)	15(25.0%)	
Clinical pregnancy			0.07
No	36(60.0%)	46(76.7%)	
Yes	24(40.0%)	14(23.3%)	




Figure 4. Endometrial thickness on the trigger day.

The comparison of clinical hirsutism between Group A and Group B (Table 3) showed a higher proportion of individuals experienced clinical hirsutism in Group A (41.7%), compared to just 16.7% in Group B.

In Table 4, clinical outcomes to ovulation induction was compared. A higher proportion of individuals in Group B (100%) had mature follicles compared to Group A (88.3%). Still, it was found that Group B (93.3%) achieved higher ovulation compared to Group A (70%), with a P-value of 0.002. In terms of pregnancy, Group A had a higher percentage of po-

sitive S.βhCG results (40.0%), compared to Group B (25.0%), and higher clinical pregnancy rate (40% vs 23.3%) but not statistically significant.

DISCUSSION

PCOS is the most prevalent endocrine disorder in the reproductive age of women. In these patients, medications such as clomiphene citrate, letrozole and metformin are used to induce ovulation and overcome infertility [12].

Clomiphene citrate binds to estrogen receptors for prolonged periods *i.e.* weeks (2 weeks) rather than hours as with natural estrogen. This extended binding ultimately depletes estrogen receptors' concentrations by interfering with the normal process estrogen receptors' replenishment and may be responsible for the peripheral antiestrogenic effect of clomiphene citrate on the endometrium and cervix [13].

Compared to CC, letrozole is cleared from circulation more rapidly due to shorter half-life; associated with monofollicular growth and thicker endometrium. Letrozole is devoid of any antiestrogenic peripheral actions and has no impact on the endometrial receptivity and cervical mucous quality [14]. Metformin, an insulin sensitizer may be used alone or in concert with other medications such as clomiphene citrate, letrozole it has been shown to increase the ovulatory response to clomiphene citrate in patients who were previously clomiphene resistant [15].

In our study, demographic parameters *i.e.* age, BMI, type of infertility was comparable in both groups. The only significant demographic character was menstrual cycle regularity. As women were selected randomly in the study, cycles were more irregular in letrozole-metformin group, but cycle regularity is not the sole criterion for metformin use [4]. A study also has reported similar age, BMI and duration of infertility distribution in both groups [16].

In Group A, patients had higher insulin PP, day 2 S.FSH, S. LH, S.E2 and S.AMH levels. The clinical hirsutism was also higher in metformin-letrozole group.

When ovulation parameters were compared, it was found that group B participants had higher achievement of formation of mature follicle than Group A, and ovulation rate was higher. In a different study no significant differences between the letrozole and the metformin-letrozole groups regarding ovulation rate, number of patients with growing

follicle, number of follicles ≥ 18 at day 12, number of ruptured follicles after 48 hours of injection of hCG, serum E2, parity, clinical presentation, period of infertility, FSH, LH were found [16].

In our study, clinical pregnancy rate was slightly higher in Group A, but not statistically significant. A retrospective study on 268 women undergoing ovulation induction/IUI in 2012-2016 has found that the addition of metformin to Letrozole does not improve follicular recruitment or pregnancy rates in OI/IUI cycles in infertile PCOS women [17]. Liu *et al.* revealed a pregnancy rate of 57.9% in letrozole in addition to metformin gathering and just 46.8% in patients who got letrozole alone [18].

In our study the endometrial thickness was comparable in both groups (8.83 vs 8.80). A study by Morad *et al.* found superior endometrial thickness with letrozole than clomiphene citrate showing CC antiestrogenic effects [19]. A significant improvement in endometrial thickness in metformin-letrozole group was found in other work [16]. On other hand, EL-Gharib *et al.* found non-significant change between endometrial thickness in both letrozole and letrozole metformin groups, like our study results [20].

The risk of gestational diabetes in pregnancy is high in PCOS especially androgenic PCOS. The hormonal and metabolic profile of hyperandrogenic PCOS patients includes insulin resistance, and an elevated androgen levels both of which are risk factors for the development of GDM. Women with PCOS are at further risk of obstetric complications including hypertensive disorders, preterm birth, induction of labour and caesarean delivery [21]. Similarly in IVF conceived pregnancies risk of GDM is high as many females have significant risk factors for GDM, such as advanced maternal age, obesity, multiple pregnancies and PCOS, suggesting a potential relationship between GDM and ART [22].

Limitations

The study involved a smaller number of participants. In clinical situations, individual treatment dosages varied, and it was not fully considered in our study. The study's limited duration prevented a thorough examination of longer-term outcomes.

CONCLUSIONS

PCOS is a complex disorder that requires a comprehensive approach to management, addressing

both reproductive and metabolic concerns. While multiple treatment options are available for PCOS, a combination of lifestyle changes, medication, and, in some cases, surgical interventions, may be necessary to address the diverse manifestations of the disorder. Only letrozole 2.5 mg BD given from day 2 for ovulation induction and performing follicular monitoring is the standard protocol in PCOS from our research, as adding metformin to letrozole does insignificant improvement in ovulation patterns in PCOS anovulatory women. The ovulation rate was slightly higher in letrozole group. Similarly, pregnancy rates were also insignificant in both groups.

COMPLIANCE WITH ETHICAL STANDARDS

Authors' contribution

S.C.: Conceptualization, writing – original draft, Writing – review & editing, data curation. R.H.: writing – review & editing. M.G.D.: Supervision.

Funding

None.

Study registration

This study was not prospectively registered and institutional ethics committee approval was obtained before participants recruitment.

Disclosure of interests

The authors declare that they have no conflict of interests.

Ethical approval

This study was approved by the Ethics Committee of Ridge IVF, Gouri Hospital (No. RIDGE/HEC/001/2025 date: 29/4/2024).

Informed consent

Informed consent was obtained from all participants.

Data sharing

Data are available under reasonable request to the corresponding author.

ACKNOWLEDGEMENTS

We would like to thank Dr. Piyush for the valuable assistance with the formal analysis.

REFERENCES

1. Diamanti-Kandarakis E, Kouli CR, Bergiele AT, Filandra FA, Tsianateli TC, Spina GG, et al. A survey of the polycystic ovary syndrome in the Greek island of Lesbos: hormonal and metabolic profile. *J Clin Endocrinol Metab.* 1999;84(11):4006-11. doi: 10.1210/jcem.84.11.6148.
2. Tehrani FR, Simbar M, Tohidi M, Hosseinpah F, Azizi F. The prevalence of polycystic ovary syndrome in a community sample of Iranian population: Iranian PCOS prevalence study. *Reprod Biol Endocrinol.* 2011;9:39. doi: 10.1186/1477-7827-9-39. P
3. Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. *Fertil Steril.* 2004;81(1):19-25. doi: 10.1016/j.fertnstert.2003.10.004.
4. Teede HJ, Tay CT, Laven JJE, Dokras A, Moran LJ, Piltonen TT, et al. Recommendations from the 2023 international evidence-based guideline for the assessment and management of polycystic ovary syndrome. *Eur J Endocrinol.* 2023;189(2):G43-G64. doi: 10.1093/ejendo/lvad096.
5. Baptiste CG, Battista MC, Trottier A, Baillargeon JP. Insulin and hyperandrogenism in women with polycystic ovary syndrome. *J Steroid Biochem Mol Biol.* 2010;122(1-3):42-52. doi: 10.1016/j.jsbmb.2009.12.010.
6. Liu X, Zhang J, Wang S. Global, regional, and national burden of infertility attributable to PCOS, 1990-2019. *Hum Reprod.* 2024;39(1):108-118. doi: 10.1093/humrep/dead241.
7. Azziz R, Carmina E, Dewailly D, Diamanti-Kandarakis E, Escobar-Morreale HF, Futterweit W, et al. Positions statement: criteria for defining polycystic ovary syndrome as a predominantly hyperandrogenic syndrome: an Androgen Excess Society guideline. *J Clin Endocrinol Metab.* 2006;91(11):4237-45. doi: 10.1210/jc.2006-0178.
8. Bailey CJ, Turner RC. Metformin. *N Engl J Med.* 1996;334(9):574-9. doi: 10.1056/NEJM199602293340906.
9. Diamanti-Kandarakis E, Christakou CD, Kandarakis E, Economou FN. Metformin: an old medication of new fashion: evolving new molecular mechanisms and clinical implications in polycystic ovary syndrome. *Eur J Endocrinol.* 2010;162(2):193-212. doi: 10.1530/EJE-09-0733.

10. Buzdar AU, Robertson JF, Eiermann W, Nabholz JM. An overview of the pharmacology and pharmacokinetics of the newer generation aromatase inhibitors anastrozole, letrozole, and exemestane. *Cancer*. 2002;95(9):2006-16. doi: 10.1002/cncr.10908.
11. El Omda F, Abd El Fattah A, Hashish M, Khalil M. A Randomized Trial of Combined Letrozole with Metformin Versus Clomiphene Citrate with Metformin in Polycystic Ovary (PCOS) patients. *Egypt J Hospital Med*. 2018;72(4):4382-4388. doi: 10.21608/ejhm.2018.9294.
12. Farshchian N, Teimuri Nezhad S, Bahrami Kamangar P. The combination of letrozole and metformin has a better therapeutic effect on uterine and ovarian arteries in PCOS patients than the combination of clomiphene citrate and metformin. *AMJ*. 2018;11(6):326-330. doi: 10.21767/AMJ.2018.3252.
13. Greenblatt RB, Barfield WE, Jungck EC, Ray AW. Induction of ovulation with MRL/41. Preliminary report. *JAMA*. 1961;178:101-4. doi: 10.1001/jama.1961.03040410001001.
14. Pritts EA. Letrozole for ovulation induction and controlled ovarian hyperstimulation. *Curr Opin Obstet Gynecol*. 2010;22(4):289-94. doi: 10.1097/GCO.0b013e32833beebf.
15. Sohrabvand F, Ansari Sh, Bagheri M. Efficacy of combined metformin-letrozole in comparison with metformin-clomiphene citrate in clomiphene-resistant infertile women with polycystic ovarian disease. *Hum Reprod*. 2006;21(6):1432-5. doi: 10.1093/humrep/del020.
16. Hamada S, Abdallah K, Samir K. Comparative Study Between the Effect of Letrozole Versus Letrozole with Metformin in Treatment of Anovulation in Overweight Women. *AIMJ*. 2022;3(7):89-95
17. Hurley EG, Adams SR, Kalakota N, Rios J, Thomas M. The addition of metformin during ovulation induction with letrozole does not affect pregnancy outcome in infertile women with polycystic ovary syndrome. *Fertil Steril*. 2017;108:e246. doi: 10.1016/j.fertnstert.2017.07.740.
18. Liu C, Feng G, Huang W, Wang Q, Yang S, Tan J, et al. Comparison of clomiphene citrate and letrozole for ovulation induction in women with polycystic ovary syndrome: a prospective randomized trial. *Gynecol Endocrinol*. 2017;33(11):872-876. doi: 10.1080/09513590.2017.1332174.
19. Morad AWA, Elhadi Farag MA. Impact of letrozole on ultrasonographic markers of endometrial receptivity in polycystic ovarysyndrome women with poor endometrial response to clomiphene citrate despite adequate ovulation. *Middle East Fertil Soc J*. 2015;20(3):182-187. doi: 10.1016/j.mefs.2014.08.004
20. EL-Gharib MN, EL-Ebiary MT, Raoof Farahat MA (2020) The Negligible Effect of Metformin Addition to Letrozole in Treating Overweight Women with PCOS. *J Obst Gynecol Surg*. 2020;1(2):1-6.
21. Quaresima P, Myers SH, Pintaudi B, D'Anna R, Morelli M, Unfer V. Gestational diabetes mellitus and polycystic ovary syndrome, a position statement from EGOI-PCOS. *Front Endocrinol (Lausanne)*. 2025;16:1501110. doi: 10.3389/fendo.2025.1501110.
22. Ghanem YM, El Kassar Y, Magdy MM, Amara M, Amin NG. Potential risk of gestational diabetes mellitus in females undergoing in vitro fertilization: a pilot study. *Clin Diabetes Endocrinol*. 2024;10(1):7. doi: 10.1186/s40842-024-00164-x.