



# Italian Journal of Gynæcology & Obstetrics

December 2024 - Vol. 36 - N. 4 - Quarterly - ISSN 2385 – 0868

## Intimate partner violence onset during wanted and unwanted pregnancy in Iranian women

Fateme Khajoei Nejad<sup>1</sup>, Katayoun Alidousti<sup>2,\*</sup>, Nahid Bagheri<sup>3</sup>

<sup>1</sup>Department of Midwifery, Sirjan School of Medical Sciences, Sirjan, Iran.

<sup>2</sup>Neuroscience Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Kerman, Iran.

<sup>3</sup>Department of Epidemiology and Biostatistics, School of Health, Kerman University of Medical Sciences, Kerman, Iran.

### ARTICLE INFO

#### History

Received: 29 March 2024

Received in revised form: 25 June 2024

Accepted: 30 July 2024

Available online: 17 December 2024

DOI: 10.36129/jog.2024.174

#### Key words

Unwanted pregnancy; maternal health; intimate partner violence.

\*Corresponding author: Katayoun Alidousti, MSc. Neuroscience Research Center, Institute of Neuropharmacology, Kerman University of Medical Sciences, Haft Bagh Alavi, Kerman, Iran. Email: alidoosti@kmu.ac.ir; kalidousti@gmail.com. ORCID: 0000-0002-1206-4912.

### ABSTRACT

**Objective.** Violence during pregnancy may lead to pregnancy complications, adverse birth outcomes, or other serious reproductive events. Unwanted pregnancy can lead to a wide range of physical and psychological consequences for both the mother and child, impacting various aspects of social, economic, and cultural health. So, this study aimed to compare the onset of intimate partner violence during wanted and unwanted pregnancies.

**Materials and Methods.** Analytical cross-sectional research was conducted using convenience sampling on 300 eligible women (150 with unwanted pregnancies and 150 with wanted pregnancies) who sought care at health care centres in Kerman in 2022. The Iranian version of the Domestic Violence Questionnaire and SPSS-22 were utilized for data collection and analysis.

**Results.** The mean age was  $29.77 \pm 7.06$  years. The majority of women (28%) had a diploma, while 69.65% were housewives. The average level of violence during unwanted pregnancy ( $60.64 \pm 30.51$ ) was higher than before pregnancy ( $51.83 \pm 25.02$ ). The mean score of domestic violence in unwanted pregnancies ( $60.64 \pm 30.51$ ) was higher than in the group with wanted pregnancies ( $51.35 \pm 26.99$ ) ( $p < 0.05$ ). The mean score of the psychological dimension in unwanted pregnancy ( $14.23 \pm 6.03$ ) and the wanted pregnancy group ( $10.57 \pm 5.32$ ) showed a statistically significant difference ( $p < 0.05$ ). Also, the mean score of patriarchal beliefs in the group with unwanted pregnancy ( $36.93 \pm 20.58$ ) during pregnancy was higher than in the group with wanted pregnancy ( $31.48 \pm 18.43$ ) ( $p < 0.05$ ).

**Conclusions.** Unwanted pregnancies can trigger the onset of domestic violence. Careful attention should be given to the topic of unwanted pregnancies when making decisions about providing training and consultation services to parents in health centres.

### INTRODUCTION

Women, especially mothers, play pivotal roles in both pedagogical and familial management affairs,

forming emotional and affective bonds among family members [1]. However, the intrusion of gender prototypes into these relations has led to patriarchal standards, indirectly contributing to domestic

violence (DV) against women. The World Health Organization recognizes DV as a global public health issue due to its prevalence and the associated physical and psychological morbidity and mortality [2]. DV, also known as intimate partner violence (IPV), includes physical, sexual, psychological, and financial aspects, along with controlling or coercive behaviours. Statistically, one in three women experiences physical or sexual violence at some point in their lives [3, 4].

Violence against women (VAW), particularly DV, has become a pervasive and challenging social issue that transcends regional and cultural boundaries in recent decades. The prevalence of physical or sexual violence against wives varies widely across countries [5]. In the United States alone, 2-4 million women face IPV each year [6]. Disturbingly, around seventeen percent of women experience DV for the first time during pregnancy [7]. DV during pregnancy is associated with various adverse outcomes, such as pregnancy complications, adverse birth outcomes, non-use of contraception, unwanted pregnancies, and increased mortality for both the mother and child [8-10]. A pregnancy accompanied by violence is considered a high-risk pregnancy. Studies have shown that in high-risk pregnancies, the likelihood of pregnancy complications and issues during the delivery process, such as induction and labour management, is higher [11, 12].

Unwanted pregnancies, whether resulting in continuation or termination, have significant physical and psychological consequences for mothers and children, affecting various aspects of social, economic, and cultural health in the community [13, 14]. Studies have shown that pregnant women exposed to DV may face traumatic outcomes, including abortion, preterm childbirth, placental separation, chorioamnionitis, abnormal bleeding of the genital system, pelvic inflammatory diseases, and increased maternal mortality [15]. In settings such as Vietnam, higher levels of DV exposure to domestic violence have been associated with reduced well-being for children [16]. Despite the recognized adverse physical and mental health consequences, DV during pregnancy remains an unresolved and neglected social problem [17].

This study aims to compare the onset of intimate partner violence during wanted and unwanted pregnancies, with the goal of contributing to the improvement of maternal and child health.

## MATERIALS AND METHODS

This analytical cross-sectional research was conducted between April and October 2022, focusing on comparing intimate partner violence among women with wanted and unwanted pregnancies; investigating whether unwanted pregnancies can lead to the initiation of intimate partner violence.

### *Sample and population*

The study population included all pregnant women receiving prenatal care at health centres in Kerman. The sample size, estimated at 300 women (150 with wanted pregnancies and 150 with unwanted pregnancies), was based on a 50% prevalence of unwanted pregnancies [18]. Random sampling was conducted across 19 clinics in Kerman, taking into account the rates of both wanted and unwanted pregnancies in each clinic.

Inclusion criteria included couples of Iranian nationality, women aged between 15 and 49 years, married, and gestational age over 20 weeks.

Exclusion criteria included drug abuse by either partner, neuropsychological disorders or diseases in either partner, experiencing stressful conditions during pregnancy (such as losing a job, moving to a new house, onset of a critical illness, loss of a friend or family member).

### *Instruments*

Data were collected through questionnaires

#### *Demographic questionnaire*

Validated by ten members of the midwifery faculty from the current study institution, this questionnaire included information on the age of the mother and father, their education, occupation, number of children, income, and length of marriage.

#### *Domestic violence questionnaire*

Developed by Mohseni Tabrizi *et al.* for the Iranian population, this questionnaire measures domestic violence using demographic information and 60 items related to physical, sexual, economic, psychological violence, and patriarchal beliefs [19]. Scoring the questionnaire is in two sections and is based on Likert's 5-point scale as follows:

Some items are responded to be based on a spectrum from "never" to "always" (*e.g.*, never, seldom, sometimes, often, and always), while other items are re-

sponded to base on a different spectrum from completely disagree to completely agree (e.g., completely disagree, disagree, no idea, agree, completely agree). Subjects can select one of the options as their response to the item. The scoring ranges from 0 to 4, where 0 represents "never" or "completely disagree," 1 represents "seldom" or "disagree," 2 represents "sometimes" or "no idea," 3 represents "agree" or "often," and 4 represents "completely agree" or "always." The minimum score is 0, and the maximum score is 240. A score between 0 and 60 indicates low levels of domestic violence against women. Score between 60 and 120: The level of domestic violence against women is considered moderate. Above 120: The rate of domestic violence against women is high.

### *Inspection of unwanted pregnancy*

Unintended pregnancy was assessed using the London Measure of Unplanned Pregnancy (LMUP) questionnaire. Six items (questions) constitute the self-administered retrospective LMUP, which assesses the level of pregnancy planning and intention for recent or ongoing pregnancies. The tool, which was initially created in the UK, has been validated in several countries and in various language versions. The topics covered include the use of contraceptives, timing of pregnancy, personal desire, influence from partners, and preparedness. For every item, a score of zero, one, or two can be obtained. An overall pregnancy intention score, which ranges from zero to twelve, is calculated by adding all of the scores together. Higher scores indicate greater levels of planning and intention. Scores should be categorized as planned if they are  $\geq 10$  and unplanned if they are 0-9 [20].

### *Procedure*

After obtaining approval from the ethics committee and written informed consent, questionnaires were distributed privately to eligible participants. Women with wanted and unwanted pregnancies were interviewed and completed the violence questionnaire twice: once reflecting on their pre-pregnancy situation retrospectively, and once regarding their current condition during pregnancy.

### *Patient and public involvement*

Before the study materials were created, three informal scoping groups were organized, each consist-

ing of 20 pregnant women. These groups engaged in discussions regarding unwanted pregnancy and partner violence, focusing on the issues that pregnant women considered significant to include. The proposed materials were then presented to these groups, fostering collaboration and enabling the pregnant women to actively contribute to the survey's content and wording. The findings of the study will be shared with the study participants and local stakeholders through educational and public health institutions.

### *Statistical analysis*

Data analysis was conducted using SPSS version 22, utilizing descriptive statistical tests to present measurement values such as mean, standard deviation, and frequency. A comparison was conducted between the average violence scores before and during pregnancy, as well as between the groups with wanted and unwanted pregnancies.

## **RESULTS**

Demographic data analysis showed that the mean age and age difference with spouse were  $29.77 \pm 7.06$  years and  $4.31 \pm 3.03$  years, respectively. 56.33% had been married for more than 6 years. 14.68% had more than 2 children. More than one third of the sample, 37.3%, had more than one child. The majority of women (28%) had a diploma, were housewives (69.65%), and had no personal income (68.31%). 40.98% of husbands had education below a diploma. 4.33% of men were unemployed, and 35.01% of men had an average income (**Table 1**).

Comparing domestic violence in the unwanted pregnancy group revealed that the overall level of violence during pregnancy ( $60.64 \pm 30.51$ ) was higher than before pregnancy ( $51.83 \pm 25.02$ ), and this difference was statistically significant ( $p < 0.05$ ). The comparison of various dimensions of violence revealed that the average score for psychological dimensions during pregnancy ( $14.23 \pm 6.03$ ) was higher than before pregnancy ( $11.08 \pm 3.22$ ), and this difference was statistically significant ( $p < 0.05$ ). The mean score of patriarchal beliefs increased from  $31.46 \pm 18.74$  before pregnancy to  $36.93 \pm 20.58$  during pregnancy, showing a significant difference ( $p < 0.05$ ). Other dimensions of domestic violence did not show significant differences between the two time points (**Table 2**).

**Table 1.** Participants' demographic characteristics

Variable		Unwanted pregnancy	Wanted pregnancy
		Mean ± SD	
Age (year)		29.89 ± 7.04	29.66 ± 7.08
Age Different with Spouse (year)		4.47 ± 3.01	4.15 ± 3.05
Variable		Frequency n (%)	
Number of children	No child	36 (24)	34 (22.66)
	1 child	58 (38.7)	56 (37.33)
	2 children	35 (23.3)	37 (24.66)
	3 children	15 (10)	16 (10.66)
	> 4 children	6 (4)	7 (4.66)
Level of education	Illiterate	2 (13)	3 (2)
	Primary	15 (10)	15 (10)
	Secondary	22 (14.7)	24 (16)
	High school	10 (6.7)	12 (8)
	Diploma	44 (29.3)	40 (26.66)
	Associate	14 (9.3)	12 (8)
	Undergraduate	37 (24.7)	36 (24)
Occupation	Graduate	6 (4)	8 (5.33)
	Employee	30 (20)	29 (19.33)
	Worker	1 (0.7)	2 (1.33)
	Self-employed	11 (7.3)	10 (6.66)
	Unemployed	4 (2.7)	4 (2.66)
	Housewife	104 (69.3)	105 (70)
Income level	No income	104 (69.3)	101 (67.33)
	< \$80	8 (5.3)	9 (6)
	\$80-160	12 (8)	11 (7.33)
	\$160-240	25 (16.7)	27 (18)
	> \$240	1 (7)	2 (1.33)
Spouse's occupation	Employee	40 (26.7)	41 (27.33)
	Worker	28 (18.7)	28 (18.66)
	Self-employed	76 (50.7)	74 (49.33)
	Unemployed	6 (4)	7 (4.66)
Year of marriage	<1 year	10 (6.7)	9 (6)
	1-3 years	35 (23.3)	36 (24)
	4-5 years	21 (14)	20 (13.33)
	> 6 years	84 (56)	85 (56.66)
Spouse's income	No income	1 (7)	2 (1.33)
	< \$80	21 (14)	19 (12.66)
	\$80-160	52 (34.7)	53 (35.33)
	\$160-240	40 (26.7)	42 (28)
	> \$240	36 (24)	34 (22.66)
Spouse's education level	Illiterate	4 (2.7)	3 (2)
	Primary	14 (9.3)	12 (8)
	Secondary	24 (16)	25 (16.66)
	High school	12 (8)	14 (9.3)
	Diploma	46 (30.7)	43 (28.66)
	Associate	11 (7.3)	10 (6.66)
	Undergraduate	27 (18)	29 (19.33)
Graduate	12 (8)	14 (9.3)	

**Table 2.** Comparison of intimate partner violence in unwanted pregnancy (before and during pregnancy).

Variable	Before pregnancy	During pregnancy	P-value
	Mean ± SD		
Sexual violence	2.15 ± 0.85	2.24 ± 0.98	0.46
Physical violence	3.98 ± 1.12	4.01 ± 1.32	0.38
Psychological violence	11.08 ± 3.22	14.23 ± 6.03	< 0.05
Economic violence	3.16 ± 1.09	3.23 ± 1.6	0.43
Patriarchic beliefs	31.46 ± 18.74	36.93 ± 20.58	< 0.05
Total score	51.83 ± 25.02	60.64 ± 30.51	< 0.05

The overall mean score of domestic violence in the group with unwanted pregnancy (60.64 ± 30.51) was higher than in the group with wanted pregnancy (51.35 ± 26.99) ( $p < 0.05$ ). By comparing the dimensions of domestic violence, we observed a statistically significant difference in the mean score of the psychological dimension between unwanted pregnancy (14.23 ± 6.03) and wanted pregnancy groups (10.57 ± 5.32) ( $p < 0.05$ ). Also, the mean score of patriarchal beliefs in the group with unwanted pregnancy (36.93 ± 20.58) during pregnancy was higher than the group with wanted pregnancy (31.48 ± 18.43) ( $p < 0.05$ ) (Table 3).

**Table 3.** Comparison of intimate partner violence during pregnancy between two groups (with wanted and unwanted pregnancy).

Variable	Unwanted pregnancy	Wanted pregnancy	P-value
	Mean ± SD		
Sexual violence	2.24 ± 0.98	2.11 ± 0.86	0.48
Physical violence	4.01 ± 1.32	4.07 ± 1.28	0.38
Psychological violence	14.23 ± 6.03	10.57 ± 5.32	< 0.05
Economic violence	3.23 ± 1.6	3.12 ± 1.1	0.41
Patriarchic beliefs	36.93 ± 20.58	31.48 ± 18.43	< 0.05
Total score	60.64 ± 30.51	51.35 ± 26.99	< 0.05

## DISCUSSION

The results of this study imply that the mean DV score among women with unwanted pregnancies is at the borderline of moderate violence. Based on Raziani's research, unwanted pregnant women experienced DV more frequently than others [21]. Other studies have also indicated that violence during pregnancy is more common when the pregnancy is unwanted [15]. Sarayloo also concluded that unwanted pregnancy is a factor in a high level of DV [22].

Given that the majority of Iranians are Muslims, based on their religious beliefs, some people consider the challenges they encounter in life as divine tests, aiming to enhance their tolerance and acceptance. In this study, we observed a shift in violent behaviours from physical actions to mental states. Men attempted to perceive unwanted pregnancy as a test from God and refrained from physically abusing women.

Investigating the dimensions of domestic violence, it was found that the majority of the subjects experienced a low level of DV. However, some researchers' results disagreed with ours and suggested that pregnant women with unwanted pregnancies were highly influenced by psychological DV followed by physical and sexual types [23]. In the current study, the psychological DV observed most frequently includes enforced isolation from friends, family, or work, swearing and cursing, verbal aggression, insults, humiliation, defamation, and threats of physical harm to oneself, partner, or children.

In a study in Uganda, it was implied that women with unwanted pregnancies were 4.1 times more likely to experience physical violence from their spouse or partner compared to women with wanted pregnancies [24]. A meta-analysis showed that the prevalence of violence among pregnant women was 17% for physical violence, 41% for psychological violence, and 21% for sexual violence. This suggests a lower level of physical violence compared to other types [25]. The transition from a traditional to a modern society involves a shift in the behavioural and discourse models that govern the society. This may justify the low level of physical violence in the current study.

The level of sexual domestic violence among the participants in this study was low. It included having sex without the wife's consent or desire, requesting anal sex against the wife's wishes, and having sex without foreplay. In a study, it was also indicated that 47.5% of women who experienced domestic violence from their spouses reported 34.2% of physical DV and 2.7% of sexual DV [26]. In contrast to the present study, a systematic study implied that sexual DV in Iranian women is moderate and more prevalent than physical violence [27].

Sexual violence consequences are multifaceted and varied, including vaginal problems, recurrent urinary tract infections, widespread and chronic pain, sleeping problems, chronic back problems, fibromyalgia, eating disorders, social anxiety, severe

depression, and chronic fatigue. In conclusion, sexual violence has extremely negative and long-term consequences due to the interconnectedness of the body, mind, and soul [28].

The level of economic violence experienced by pregnant women was low. Aghakhani found that 88% of women experience economic violence, in contrast to this study [29]. Talebpour also reported that the highest level of violence is associated with the financial dimension [30]. The geographical and age differences, along with customs and traditions of a society and ethnicity, can be the reasons for different results.

The results found that the level of violence was moderate in the dimension of patriarchal beliefs. One of the sources of marital conflict is patriarchal attitudes. Patriarchy refers to a system that socially, politically, and economically dominated women. The patriarchal orientations and attitudes directly influence violence against women. A study in India reported that among various reasons for high prevalence of domestic violence, the patriarchal roles and long-term cultural norms that subordinate women are the most profound forms of violence [31]. Zare Shahabadi concluded that patriarchy is directly related to psychological, physical, and financial DV [32]. A sociological study revealed that the intervention of the husband's family and friends in their family's issues, making decisions on their behalf, the patriarchal behaviour of the husband (influenced by his upbringing environment), the presence of violence within the husband's family, and the woman's secondary role in the husband's life were identified as factors contributing to violent behaviour [33].

However, Iranian society is progressing towards modernity, and physical violence is becoming less prevalent. Nevertheless, patriarchal beliefs still persist in society, and this mindset can be altered through long-term training and cultural interventions.

In Iran, the Civil Code, Islamic Penal Code, Development Program Laws, Regulations, Legal Bills, and Charters of Citizens' Rights have prioritized the protection of women and the fight against violence targeting women. However, most of these laws have not specified their implementation method and scope after approval [34]. It is necessary to amend the previous laws and establish new laws in accordance with the current needs of the society and international standards, as seen in some countries [35].

## CONCLUSIONS

The study indicates a higher rate of intimate partner violence against women with unwanted pregnancies and suggests that such pregnancies can trigger the onset of domestic violence. Given the unique mental and physical conditions of pregnancy, unwanted pregnancies exacerbate these conditions, leading to additional stress. Interventions aimed at supporting women with unwanted pregnancies can contribute to reducing maternal and neonatal mortality and morbidity, promoting overall health.

### Limitations

The study had limitations, including data availability only for women with a gestational age over 20 weeks. This limited the assessment of the impact of intimate partner violence (IPV) on unwanted pregnancy for those who experienced miscarriage or induced abortion. Additionally, reliance on self-reported data may lead to underreporting of domestic violence due to societal or personal reasons.

## COMPLIANCE WITH ETHICAL STANDARDS

### Authors' contribution

F.K.H., K.A., N.B.: Conceptualization. F.K.H., N.B., K.A.: Data curation, formal analysis. K.A.: writing – original draft, writing – review & editing.

### Funding

None.

### Study registration

N/A.

### Disclosure of interests

The authors declare that they have no conflict of interests.

### Ethical approval

The study was approved by the Ethics Committee of Kerman University of Medical Sciences, Iran

(Ethics code: IR.KMU.REC.1399.125), and adhered to the Declaration of Helsinki and COPE guidelines.

### Informed consent

Written informed consent was obtained, and participants had the option to withdraw from the study at any time.

### Data sharing

Data are available under reasonable request to the corresponding author.

## REFERENCES

1. Muluneh MD, Francis L, Agho K, Stulz V. The association of intimate partner violence and contraceptive use: a multi-country analysis of demographic and health surveys. *Int J Equity Health*. 2023;22(1):75. doi: 10.1186/s12939-023-01884-9.
2. Bazargan-Hejazi S, Medeiros S, Mohammadi R, Lin J, Dalal K. Patterns of intimate partner violence: a study of female victims in Malawi. *Journal of Injury and Violence Research*. 2013;5(1):38. doi: 10.5249/jivr.v5i1.139.
3. Devries KM, Mak JY, Bacchus LJ, Child JC, Falder G, Petzold M, et al. Intimate partner violence and incident depressive symptoms and suicide attempts: a systematic review of longitudinal studies. *PLoS Med*. 2013;10(5):e1001439. doi: 10.1371/journal.pmed.1001439.
4. Bacchus LJ, Ranganathan M, Watts C, Devries K. Recent intimate partner violence against women and health: a systematic review and meta-analysis of cohort studies. *BMJ open*. 2018;8(7):e019995. doi: 10.1136/bmjopen-2017-019995.
5. Bola SL. Spousal violence and pregnancy termination among married women in Nigeria. *Afr Health Sci*. 2016;16(2):429-40. doi:10.4314/ahs.v16i2.11.
6. Tiyyagura G, Christian C, Berger R, Lindberg D; ExSTRA Investigators. Occult abusive injuries in children brought for care after intimate partner violence: An exploratory study. *Child Abuse Negl*. 2018;79:136-43. doi: 10.1016/j.chiabu.2018.02.003.
7. Baird K, Creed DK, Saito AS, Eustace J. Longitudinal evaluation of a training program to promote

- routine antenatal enquiry for domestic violence by midwives. *Women Birth*. 2018;31(5):398-406. doi: 10.1016/j.wombi.2018.01.004.
8. Felker-Kantor E, Wallace M, Theall K. Living in violence: Neighborhood domestic violence and small for gestational age births. *Health Place*. 2017;46:130-6. doi: 10.1016/j.healthplace.2017.05.011.
  9. Fite RO, Mohammedamin A, Abebe TW. Unintended pregnancy and associated factors among pregnant women in Arsi Negele Woreda, West Arsi Zone, Ethiopia. *BMC Research Notes*. 2018;11(1):1-7. doi: 10.1186/s13104-018-3778-7.
  10. Faraidoon R, Ali SM. Birth weight in high risk and normal pregnancy: a comparative study. *Ital J Gynaecol Obstet*. 2024;36(4):455-65. doi: 10.36129/jog.2024.164.
  11. Sfregola G, Sfregola P, Ruta F, Zendoli F, Musicco A, Garzon S, et al. Effect of maternal age and body mass index on induction of labor with oral misoprostol for premature rupture of membrane at term: A retrospective cross-sectional study. *Open Med*. 2023;18(1):20230747. doi: 10.1515/med-2023-0747.
  12. Etrusco A, Sfregola G, Zendoli F, Musicco A, Belpiede A, Della Pietà C, et al. Effect of maternal age and BMI on induction of labor using oral misoprostol in late-term pregnancies: a retrospective cross-sectional study. *Gynecol Obstet Invest*. 2024;89(4):323-9. doi: 10.1159/000538374.
  13. Bishwajit G, Tang S, Yaya S, Feng Z. Unmet need for contraception and its association with unintended pregnancy in Bangladesh. *BMC pregnancy and childbirth*. 2017;17(1):1-9. doi: 10.1186/s12884-017-1379-4.
  14. Rottenstreich M, Loitner L, Dar S, Kedem R, Smorgick N, Vaknin Z. Unintended pregnancies among women serving in the Israeli military. *Contraception*. 2017;96(1):62-5. doi: 10.1016/j.contraception.2017.03.006.
  15. Ahinkorah BO, Seidu A-A, Appiah F, Oduro JK, Sambah F, Baatiema L, et al. Effect of sexual violence on wanted, mistimed and unwanted pregnancies among women of reproductive age in sub-Saharan Africa: A multi-country analysis of Demographic and Health Surveys. *SSM-population health*. 2020;11:100601. doi: 10.1016/j.ssmph.2020.100601.
  16. Bui QN, Hoang TX, Le NT. The effect of domestic violence against women on child welfare in Vietnam. *Children and Youth Services Review*. 2018;94:709-19. doi: 10.1016/j.childyouth.2018.09.024.
  17. Da Thi Tran T, Murray L, Van Vo T. Intimate partner violence during pregnancy and maternal and child health outcomes: a scoping review of the literature from low-and-middle income countries from 2016 - 2021. *BMC Pregnancy Childbirth* 2022;22(1):315. doi: 10.1186/s12884-022-04604-3.
  18. Esfahani P, Danshi Kohani Z, Arefi M. Prevalence of Unwanted Pregnancy among Iranian Women: Systematic Review and Meta - analysis. *Pajouhan Sci J*. 2020;18(2):1-12. doi: 10.52547/psj.18.2.1.
  19. Mohseni Tabrizi A, Kaldi A, Javadianzadeh M. The Study of Domestic Violence in Married Women Admitted to Yazd Legal Medicine Organization and Welfare Organization. *Tolooebehdasht*. 2013;11(3):11-24.
  20. Lang AY, Hall JA, Boyle JA, Harrison CL, Teede H, Moran LJ, Barrett G. Validation of the London Measure of Unplanned Pregnancy among pregnant Australian women. *PLoS One*. 2019;14(8):e0220774. doi: 10.1371/journal.pone.0220774.
  21. Raziani Y, Hasheminasab L, Gheshlagh RG, Dalvand P, Baghi V, Aslani M. The prevalence of intimate partner violence among Iranian pregnant women: a systematic review and meta-analysis. *Scand J Public Health*. 2024;52(1):108-18. doi: 10.1177/14034948221119641.
  22. Sarayloo K, Mirzaei Najmabadi K, Ranjbar F, Behboodi Moghadam Z. Prevalence and risk factors for domestic violence against pregnant women. *Iran Journal of Nursing*. 2017;29(104):28-35. doi: 10.29252/ijn.29.104.28.
  23. Zheng B, Zhu X, Hu Z, Zhou W, Yu Y, Yin S, et al. The prevalence of domestic violence and its association with family factors: a cross-sectional study among pregnant women in urban communities of Hengyang City, China. *BMC Public Health*. 2020;20(1):620. doi: 10.1186/s12889-020-08683-9.
  24. Hailu HT, Mekonnen W, Gufue ZH, Weldegebriel SG and Dessalegn B. Intimate partner violence as a determinant factor for spontaneous abortion during pregnancy: an unmatched case-control study. *Front. Public Health*. 2023;11:1114661. doi: 10.3389/fpubh.2023.1114661.
  25. Akhtari Zavare M, Ghaleiha A, Matinnia N. Prevalence and Risk Factors of Domestic Violence in Primigravidae in Low Socio-Economic Areas of Hamedan, Iran. *Avicenna J Neuro Psycho Physiol*. 2022;9(4):137-43- doi: 10.32592/ajnp.2022.9.4.100.

26. Potter LC, Morris R, Hegarty K, García-Moreno C, Feder G. Categories and health impacts of intimate partner violence in the World Health Organization multi-country study on women's health and domestic violence. *Int J Epidemiol.* 2021;50(2):652-62. doi: 10.1093/ije/dyaa220.
27. Zare E, Ghaffari M, Nahidi F, Nasiri M, Masjedi A. Relationship Between Domestic Violence in Pregnancy and Maternal Fetus Attachment. *Iran J Psychiatry Behav Sci.* 2022;16(1):e111406. doi: 10.5812/ijpbs.111406.
28. Sigurdardottir S, Halldorsdottir S. Persistent Suffering: The Serious Consequences of Sexual Violence against Women and Girls, Their Search for Inner Healing and the Significance of the #Me-Too Movement. *Int J Environ Res Public Health.* 2021;18(4):1849. doi: 10.3390/ijerph18041849.
29. Aghakhani N, Mosavi E, Eftekhari A, Eghtedar S, Zareei A, Rahbar N, et al. A study on the domestic violence in women with addicted and non-addicted husbands referred to forensic center of Urmia, Iran. *Journal Urmia Nurs Midw Faculty.* 2014;11(11).
30. Talebpour A. Investigating Affecting Factors on Violence against Women with Emphases on Wife Abuse (Case Study; Ardebil Province). *Two Quart J Cont Sociol Res.* 2018;6(11):27-49. doi: 10.22084/csr.2017.11789.1191.
31. Kalokhe A, Del Rio C, Dunkle K, Stephenson R, Metheny N, Paranjape A, et al. Domestic violence against women in India: A systematic review of a decade of quantitative studies. *Global public health.* 2017;12(4):498-513. doi: 10.1080/17441692.2015.1119293.
32. Zare Shahabadi A, Nadarpoor Y. An investigation into the relationship between patriarchy and violence against women in koohdasht. *Social Welfare Quart.* 2016;16(60):61-86.
33. Rahbar Taramsari M, Badsar A, Zobde Imanabadi R, Khajeh Jahromi S, Amir Maafi A, Yaghubi M. Evaluation of physical intimate partner violence in respective victims in Rasht. *J Guilan Uni Med Sci.* 2012;21(83):21-6.
34. Mahmoudi A, Majidi H. Controlling, preventing and combating violence against women in Iranian laws. *J Women Fam Stud.* 2022;56:107-28.
35. Aquino CI, Marisei M, Surico D, DI Donna G. "Short Epic" on gender-based violence in the healthcare landscape, according to Italian legislation: an example of modern social and cultural evolution. *Ital J Gynaecol Obstet.* 2024;36(1):129-31. doi: 10.36129/jog.2023.126.